



Queensland

Vegetation Management Amendment Regulation (No. 2) 2013

Subordinate Legislation 2013 No. 256

made under the

Vegetation Management Act 1999

Contents

		Page
1	Short title	2
2	Commencement.....	2
3	Regulation amended	2
4	Replacement of pt 2 (Approvals of particular policies, codes and maps)	2
	Part 2 Approval of self-assessable codes	
3	3 Approval of self-assessable vegetation clearing codes—Act, s 19P	2
5	Amendment of s 11 (Matters prescribed for property vegetation management plan)	3
6	Amendment of s 12 (Fees).....	5
7	Replacement of schs 1–3.....	5
	Schedule 1 Endangered regional ecosystems	
	Schedule 2 Of concern regional ecosystems	
	Schedule 3 Least concern regional ecosystems	
8	Amendment of sch 4 (Grassland regional ecosystems—Act, schedule)	137
9	Amendment of sch 7 (Fees).....	137

[s 1]

1 Short title

This regulation may be cited as the *Vegetation Management Amendment Regulation (No. 2) 2013*.

2 Commencement

This regulation commences on 2 December 2013.

3 Regulation amended

This regulation amends the *Vegetation Management Regulation 2012*.

4 Replacement of pt 2 (Approvals of particular policies, codes and maps)

Part 2—

omit, insert—

**Part 2 Approval of
 self-assessable codes**

3 Approval of self-assessable vegetation clearing codes—Act, s 19P

The following codes are approved as self-assessable vegetation clearing codes—

- (a) the code called ‘Managing category C regrowth vegetation’ made by the Minister on 14 November 2013;
- (b) the code called ‘Managing category R regrowth vegetation’ made by the Minister on 14 November 2013;
- (c) the code called ‘Managing clearing to improve operational efficiency of existing agriculture’ made by the Minister on 14 November 2013;

[s 5]

- (d) the code called ‘Managing clearing for necessary property infrastructure’ made by the Minister on 14 November 2013;
- (e) the code called ‘Managing encroachment’ made by the Minister on 14 November 2013;
- (f) the code called ‘Managing fodder harvesting’ made by the Minister on 14 November 2013;
- (g) the code called ‘Managing native forest practice’ made by the Minister on 14 November 2013.
- (h) the code called ‘Managing thickened vegetation in the Mulga Lands’ made by the Minister on 14 November 2013;
- (i) the code called ‘Managing weeds’ made by the Minister on 14 November 2013;

Note—

The approval of codes under this section took effect when this section commenced on 2 December 2013.

5 Amendment of s 11 (Matters prescribed for property vegetation management plan)

- (1) Section 11(2)(c)—

omit, insert—

- (c) details of the way the proposed clearing achieves the performance outcomes under—
 - (i) module 8 (native vegetation clearing) of the State development assessment provisions; or
 - (ii) if the proposed clearing is within a declared area and a declared area code exists for the area—the declared area code for the area; and

[s 5]

- (d) if an offset is proposed to satisfy a performance outcome mentioned in paragraph (c)—details on how the clearing of vegetation has been avoided or minimised.
- (2) Section 11(3)(b), note—
omit.
- (3) Section 11(3)(d)—
omit, insert—
 - (d) details of the way the proposed clearing achieves the performance outcomes under module 8 (native vegetation clearing) of the State development assessment provisions; and
 - (e) if an offset is proposed to satisfy a performance outcome mentioned in paragraph (d)—details on how the clearing of vegetation has been avoided or minimised.
- (4) Section 11(4)—
omit.
- (5) Section 11(7), definitions *Cape York Peninsula Region* and *relevant regional vegetation management code*—
omit.
- (6) Section 11(7)—
insert—

State development assessment provisions see the *Sustainable Planning Regulation 2009*, schedule 26.
- (7) Section 11(5) to (7)—
renumber as section 11(4) to (6).

6 Amendment of s 12 (Fees)

Section 12, from ‘the Act’—

omit, insert—

the Act are stated in schedule 7.

7 Replacement of schs 1–3

Schedules 1 to 3—

omit, insert—

Schedule 1 Endangered regional ecosystems

section 8(1) and (6)

Part 1 Brigalow Belt Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> open forest on alluvial plains	11.3.1
Semi-evergreen vine thicket on alluvial plains	11.3.11
<i>Dichanthium sericeum</i> and/or <i>Astrebla</i> spp. grassland on alluvial plains. Cracking clay soils	11.3.21
<i>Themeda avenacea</i> grassland on alluvial plains. Basalt derived soils	11.3.24

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tereticornis, Melaleuca viridiflora, Corymbia tessellaris and Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> tall woodland with a grassy ground layer on alluvial plains and broad drainage lines derived from serpentinite	11.3.38
Semi-evergreen vine thicket ± <i>Casuarina cristata</i> on Cainozoic clay plains	11.4.1
<i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> shrubby open forest on Cainozoic clay plains	11.4.3
<i>Eucalyptus populnea</i> with <i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> open forest to woodland on Cainozoic clay plains	11.4.7
<i>Eucalyptus cambageana</i> woodland to open forest with <i>Acacia harpophylla</i> or <i>A. argyrodendron</i> on Cainozoic clay plains	11.4.8
<i>Acacia harpophylla</i> shrubby woodland with <i>Terminalia oblongata</i> on Cainozoic clay plains	11.4.9
<i>Eucalyptus populnea</i> or <i>E. woollsiana</i> , <i>Acacia harpophylla</i> , <i>Casuarina cristata</i> open forest to woodland on margins of Cainozoic clay plains	11.4.10
<i>Eucalyptus populnea</i> woodland on Cainozoic clay plains	11.4.12
<i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> open forest in depressions on Cainozoic sand plains and/or remnant surfaces	11.5.16
<i>Eucalyptus tereticornis</i> woodland in depressions on Cainozoic sand plains and remnant surfaces	11.5.17
Semi-evergreen vine thicket and microphyll vine forest on Cainozoic igneous rocks	11.8.13

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus brownii</i> or <i>Eucalyptus populnea</i> woodland on Cainozoic igneous rocks	11.8.15
<i>Acacia harpophylla-Eucalyptus cambageana</i> woodland to open forest on fine-grained sedimentary rocks	11.9.1
<i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> open forest on fine-grained sedimentary rocks	11.9.5
<i>Acacia melvillei</i> ± <i>A. harpophylla</i> open forest on fine-grained sedimentary rocks	11.9.6
<i>Dichanthium sericeum</i> grassland with clumps of <i>Acacia harpophylla</i> on fine-grained sedimentary rocks	11.9.12
<i>Acacia harpophylla</i> open forest on deformed and metamorphosed sediments and interbedded volcanics	11.11.14
Semi-evergreen vine thicket on old sedimentary rocks with varying degrees of metamorphism and folding	11.11.18
<i>Eucalyptus populnea</i> woodland on igneous rocks. Colluvial lower slopes	11.12.17
<i>Acacia harpophylla</i> open forest on igneous rocks. Colluvial lower slopes	11.12.21

[s 7]

Part 2

Cape York Peninsula Bioregion

Column 1

Regional ecosystem

Column 2

Regional ecosystem number

Complex mesophyll vine forest on basalt lowlands

3.8.1

Part 3

Central Queensland Coast Bioregion

Column 1

Regional ecosystem

Column 2

Regional ecosystem number

Melaleuca spp. and/or *Corymbia* spp. and/or

8.2.13

Lophostemon suaveolens and/or *Acacia* spp. open forest on dune sands mixed with alluvial material ± marine sediments

Melaleuca viridiflora woodland on seasonally inundated alluvial plains with impeded drainage

8.3.2

Freshwater wetlands with permanent water and aquatic vegetation

8.3.4

Melaleuca viridiflora var. *attenuata* open forest in broad drainage areas

8.3.11

Imperata cylindrica and/or *Sorghum nitidum* forma *aristatum* and/or *Ischaemum australe* tussock grassland on alluvial and old marine plains

8.3.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia clarksoniana</i> and/or <i>C. intermedia</i> open forest on Tertiary sand plains and rises including small areas of shale (mainly subregion 6)	8.5.1
<i>Melaleuca viridiflora</i> ± <i>Allocasuarina luehmannii</i> , or <i>M. viridiflora</i> and <i>M. nervosa</i> woodland, on Tertiary sand plains	8.5.2
<i>Eucalyptus platyphylla</i> and/or <i>Corymbia clarksoniana</i> and/or <i>C. intermedia</i> and/or <i>C. tessellaris</i> woodland on low undulating areas on metamorphosed sediments	8.11.4
<i>Corymbia tessellaris</i> and/or <i>Eucalyptus tereticornis</i> ± <i>C. intermedia</i> ± <i>C. clarksoniana</i> open forest with a secondary tree layer of <i>Livistona decora</i> on low hills on Mesozoic to Proterozoic igneous rocks	8.12.27

Part 4 Desert Uplands Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia cambagei</i> woodland on lakeside dunes	10.3.19

[s 7]

Part 5 Gulf Plains Bioregion

Column 1**Regional ecosystem****Column 2****Regional
ecosystem
number**

Springs on recent alluvium

2.3.39

Part 6 Mitchell Grass Downs Bioregion

Column 1**Regional ecosystem****Column 2****Regional
ecosystem
number**

Springs on recent alluvia and fine-grained sedimentary
rock

4.3.22

Part 7 Mulga Lands Bioregion

Column 1**Regional ecosystem****Column 2****Regional
ecosystem
number**

Springs on recent alluvia, ancient alluvia and
fine-grained sedimentary rock

6.3.23

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia cambagei</i> ± <i>Casuarina cristata</i> low open forest on clay plains	6.4.1
<i>Casuarina cristata</i> ± <i>Acacia harpophylla</i> open forest on clay plains	6.4.2

Part 8 New England Tableland Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus blakelyi</i> woodland on alluvial plains	13.3.1
<i>Eucalyptus nova-anglica</i> open forest on alluvial plains	13.3.2
<i>Eucalyptus nobilis</i> open forest on alluvial plains	13.3.3
<i>Eucalyptus conica</i> , <i>E. microcarpa</i> , <i>E. melliodora</i> woodland on alluvial plains	13.3.4
<i>Eucalyptus tereticornis</i> , <i>Angophora floribunda</i> open forest on alluvial plains	13.3.7
<i>Eucalyptus moluccana</i> open forest on fine-grained sedimentary rocks	13.9.2
<i>Eucalyptus melliodora</i> and/or <i>E. moluccana</i> and/or <i>E. microcarpa</i> and/or <i>E. conica</i> woodland on igneous rocks	13.12.8

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus blakelyi</i> and/or <i>E. caliginosa</i> woodland to open forest on igneous rocks	13.12.9
<i>Eucalyptus crebra</i> , <i>E. tereticornis</i> , <i>Angophora leiocarpa</i> woodland on igneous rocks	13.12.10

Part 9 South East Queensland Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Gallery rainforest (notophyll vine forest) on alluvial plains	12.3.1
<i>Eucalyptus tereticornis</i> woodland on Quaternary alluvium	12.3.3
<i>Eucalyptus populnea</i> woodland on alluvial plains	12.3.10
<i>Corymbia intermedia</i> , <i>Eucalyptus tereticornis</i> open forest on remnant Tertiary surfaces, usually near coast and in deep red soils	12.5.2
<i>Eucalyptus racemosa</i> woodland on remnant Tertiary surfaces	12.5.3
<i>Eucalyptus siderophloia</i> , <i>E. propinqua</i> , <i>E. microcorys</i> and/or <i>E. pilularis</i> open forest on remnant Tertiary surfaces, usually deep red soils	12.5.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Syncarpia glomulifera</i> woodland on complex of remnant Tertiary surface and Tertiary sedimentary rocks	12.5.11
Microphyll to notophyll vine forest ± <i>Araucaria cunninghamii</i> on remnant Tertiary surfaces	12.5.13
Semi-evergreen vine thicket with <i>Brachychiton rupestris</i> on Cainozoic igneous rocks, usually in southern half of bioregion	12.8.21
Semi-evergreen vine thicket with <i>Brachychiton australis</i> on Cainozoic igneous rocks, usually in northern half of bioregion	12.8.22
<i>Acacia harpophylla</i> open forest on Cainozoic igneous rocks	12.8.23
<i>Corymbia citriodora</i> subsp. <i>variegata</i> open forest on Cainozoic igneous rocks especially trachyte	12.8.24
<i>Dichanthium</i> spp. and <i>Themeda triandra</i> grassland on igneous rocks	12.8.27
<i>Acacia harpophylla</i> open forest on sedimentary rocks	12.9–10.6
<i>Eucalyptus melanophloia</i> , <i>E. crebra</i> woodland on sedimentary rocks	12.9–10.8
<i>Melaleuca irbyana</i> low open forest on sedimentary rocks	12.9–10.11
<i>Eucalyptus seeana</i> , <i>Corymbia intermedia</i> , <i>Angophora leiocarpa</i> woodland on sedimentary rocks	12.9–10.12
Semi-evergreen vine thicket with <i>Brachychiton rupestris</i> on sedimentary rocks	12.9–10.15

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus cloeziana</i> open forest on metamorphics ± interbedded volcanics	12.11.16
<i>Eucalyptus pilularis</i> open forest on coastal metamorphics and interbedded volcanics	12.11.23
Semi-evergreen vine thicket on Mesozoic to Proterozoic igneous rocks, usually in southern half of bioregion	12.12.17
<i>Acacia harpophylla</i> open forest on Mesozoic to Proterozoic igneous rocks	12.12.26

Part 10 Wet Tropics Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Mesophyll vine forest on beach ridges and sand plains of beach origin	7.2.1
<i>Hemarthria uncinata</i> and/or <i>Ischaemum australe</i> ± <i>Sorghum</i> spp. grassland and/or ephemeral sedgelands on seasonally inundated alluvial plains	7.3.1
<i>Melaleuca dealbata</i> ± <i>Melaleuca leucadendra</i> open forest on poorly drained alluvial plains	7.3.6
<i>Eucalyptus pellita</i> and <i>Corymbia intermedia</i> open forest to woodland (or vine forest with emergent <i>E. pellita</i> and <i>C. intermedia</i>) on poorly drained alluvial plains	7.3.7

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia tessellaris</i> , <i>Acacia</i> spp., <i>Melaleuca</i> spp. open forest on poorly drained alluvial plains (some soils with marine plain and dune influence)	7.3.9
Mixed eucalypt open forest to woodland dominated by <i>Eucalyptus tereticornis</i> and <i>Corymbia tessellaris</i> ± <i>Melaleuca dealbata</i> (or vine forest with these species as emergents). Lowland alluvial plains	7.3.12
Complex mesophyll vine forest on well-drained alluvium of high fertility	7.3.17
Simple to complex semi-deciduous notophyll to mesophyll vine forest on lowland alluvium, predominantly riverine levees	7.3.23
Complex of fernlands and sedgelands with emergent rainforest pioneering spp. in permanently wet peat swamps of alluvial plains	7.3.30
<i>Imperata cylindrica</i> and/or <i>Sorghum nitidum</i> and/or <i>Mnesithea rottboellioides</i> and/or <i>Themeda triandra</i> closed-tussock grassland on alluvial plains	7.3.32
<i>Melaleuca viridiflora</i> var. <i>attenuata</i> open forest to closed forest on broad swampy drainage lines of alluvial plains	7.3.34
<i>Acacia mangium</i> and/or <i>A. celsa</i> and/or <i>A. polystachya</i> closed forest on alluvial plains	7.3.35
Complex semi-evergreen notophyll vine forest of uplands on alluvium	7.3.37
<i>Eucalyptus tereticornis</i> open forest on well-drained alluvial plains of lowlands	7.3.40

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus leptophleba, Corymbia clarksoniana</i> open forest to woodland on alluvium in near-coastal areas with moderate rainfall	7.3.44
<i>Lophostemon suaveolens</i> open forest to woodland on alluvial plains	7.3.46
Complex semi-evergreen notophyll vine forest of uplands on basalt	7.8.3
<i>Corymbia clarksoniana</i> open forest to woodland on basalt	7.8.19

Schedule 2 Of concern regional ecosystems

section 8(2) and (6)

Part 1 Brigalow Belt Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Sedgelands on marine clay plains	11.1.3
<i>Corymbia tessellaris</i> woodland on flat coastal dunes	11.2.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Complex of <i>Spinifex sericeus</i> , <i>Ipomoea pes-caprae</i> and <i>Casuarina equisetifolia</i> grassland and hermland on fore dunes	11.2.2
Microphyll vine forest (“beach scrub”) on sandy beach ridges and dune swales	11.2.3
Lagoons in coastal dune swales	11.2.4
<i>Eucalyptus populnea</i> woodland on alluvial plains	11.3.2
<i>Eucalyptus coolabah</i> woodland on alluvial plains	11.3.3
<i>Eucalyptus tereticornis</i> and/or <i>Eucalyptus</i> spp. woodland on alluvial plains	11.3.4
<i>Grevillea striata</i> on coastal alluvial plains	11.3.13
<i>Eucalyptus coolabah</i> , <i>Acacia stenophylla</i> , <i>Muehlenbeckia florulenta</i> fringing woodland on alluvial plains	11.3.15
<i>Eucalyptus populnea</i> woodland with <i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> on alluvial plains	11.3.17
Springs associated with recent alluvia, but also including those on fine-grained sedimentary rocks, basalt, ancient alluvia and metamorphic rocks	11.3.22
<i>Eucalyptus conica</i> , <i>E. nobilis</i> , <i>E. tereticornis</i> , <i>Angophora floribunda</i> on alluvial plains. Basalt derived soils	11.3.23
<i>Eucalyptus coolabah</i> ± <i>Casuarina cristata</i> open woodland on alluvial plains	11.3.28
<i>Eremophila mitchellii</i> open woodland on alluvial plains	11.3.33

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia tephrina</i> woodland on alluvial plains	11.3.34
<i>Eucalyptus crebra</i> and/or <i>E. populnea</i> and/or <i>E. melanophloia</i> on alluvial plains. Higher terraces	11.3.36
<i>Eucalyptus</i> spp. and/or <i>Corymbia</i> spp. grassy or shrubby woodland on Cainozoic clay plains	11.4.2
<i>Acacia argyrodendron</i> woodland on Cainozoic clay plains	11.4.5
<i>Acacia cambagei</i> woodland on Cainozoic clay plains	11.4.6
<i>Dichanthium sericeum</i> , <i>Astrebla</i> spp. and patchy <i>Acacia harpophylla</i> and <i>Eucalyptus coolabah</i> on Cainozoic clay plains	11.4.11
<i>Triodia</i> spp. grassland on Cainozoic sand plains and/or remnant surfaces	11.5.6
<i>Melaleuca tamariscina</i> shrubland on Cainozoic sand plains and/or remnant surfaces	11.5.10
<i>Acacia leptostachya</i> shrubland on Cainozoic sand plains and/or remnant surfaces	11.5.11
<i>Eucalyptus populnea</i> ± <i>Acacia aneura</i> ± <i>E. melanophloia</i> woodland on Cainozoic sand plains and/or remnant surfaces	11.5.13
<i>Triodia</i> sp. grassland with emergent trees on Cainozoic sand plains and/or remnant surfaces. Highly alkaline soils	11.5.14
<i>Micromyrtus capricornia</i> shrubland on Cainozoic sand plains and/or remnant surfaces	11.5.18
Semi-evergreen vine thicket on Cainozoic igneous rocks	11.8.3

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Shrubland (heath) on Cainozoic igneous rocks	11.8.7
<i>Callitris</i> spp. ± vine thicket on Cainozoic igneous rocks	11.8.9
<i>Themeda triandra</i> grassland on Cainozoic igneous rocks	11.8.10
<i>Dichanthium sericeum</i> grassland on Cainozoic igneous rocks	11.8.11
<i>Eucalyptus microcarpa</i> , <i>E. exserta</i> woodland on Cainozoic igneous rocks	11.8.12
<i>Eucalyptus crebra</i> , <i>Corymbia dallachiana</i> woodland on Cainozoic igneous rocks	11.8.14
Semi-evergreen vine thicket or <i>Acacia harpophylla</i> with a semi-evergreen vine thicket understorey on fine-grained sedimentary rocks	11.9.4
<i>Eucalyptus populnea</i> , <i>Eremophila mitchellii</i> shrubby woodland on fine-grained sedimentary rocks	11.9.7
<i>Eucalyptus populnea</i> and/or <i>Acacia harpophylla</i> open forest on fine-grained sedimentary rocks	11.9.10
<i>Acacia harpophylla</i> shrubland on fine-grained sedimentary rocks	11.9.11
<i>Eucalyptus moluccana</i> or <i>E. microcarpa</i> open forest on fine-grained sedimentary rocks	11.9.13
<i>Lysiphylgium carronii</i> , <i>Atalaya hemiglaucha</i> ± <i>Eucalyptus melanophloia</i> ± <i>Acacia excelsa</i> open woodland	11.9.14
Tall open forest in sheltered gorges on coarse-grained sedimentary rocks	11.10.2

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Semi-evergreen vine thicket in sheltered habitats on medium to coarse-grained sedimentary rocks	11.10.8
Springs associated with sandstone	11.10.14
<i>Eucalyptus melanophloia</i> woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.10
<i>Eucalyptus orgadophila</i> woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.11
<i>Acacia harpophylla</i> or <i>A. argyrodendron</i> and/or <i>Terminalia oblongata</i> low open forest on deformed and metamorphosed sediments and interbedded volcanics	11.11.13
<i>Eucalyptus cambageana</i> , <i>Acacia harpophylla</i> woodland on old sedimentary rocks with varying degrees of metamorphism and folding. Lowlands	11.11.16
<i>Dichanthium sericeum</i> grassland on old sedimentary rocks with varying degrees of metamorphism and folding	11.11.17
Semi-evergreen vine thicket on serpentinite	11.11.21
<i>Corymbia</i> spp., <i>Lysicarpus angustifolius</i> , <i>Eucalyptus crebra</i> , <i>E. cloeziana</i> woodland on igneous rocks (granite)	11.12.5
<i>Eucalyptus shirleyi</i> woodland on igneous rocks	11.12.8
<i>Corymbia clarksoniana</i> woodland on igneous rocks	11.12.10
<i>Melaleuca</i> spp. woodland on igneous rocks. Lowlands	11.12.11
<i>Araucaria cunninghamii</i> woodland on igneous rocks. Coastal hills	11.12.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Lophostemon</i> spp. woodland on igneous rocks. Coastal hills	11.12.14
<i>Allocasuarina torulosa, Livistona decora</i> woodland on igneous rocks. Coastal hills	11.12.15
Mixed low woodland to shrubland on igneous rocks. Coastal hills	11.12.16
Montane shrubland on igneous rocks	11.12.18
<i>Eucalyptus exserta, E. moluccana, E. crebra, Corymbia citriodora</i> woodland on igneous rocks	11.12.19
<i>Corymbia</i> spp., <i>Eucalyptus baileyana, E. dura, E. exserta</i> woodland on igneous rocks	11.12.20

Part 2 **Cape York Peninsula Bioregion**

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Excoecaria agallocha ± Aegiceras corniculatum</i> closed scrub on upper tidal reaches of rivers	3.1.4
<i>Schoenoplectus</i> spp. sedgelands in depressions on tidal flats	3.1.7
Evergreen notophyll vine forest in coastal dunefield systems	3.2.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca dealbata</i> ± <i>Acacia crassarpa</i> open forest in dune swales on the west coast	3.2.3
<i>Melaleuca leucadendra</i> ± <i>M. dealbata</i> open forest in dune swales and swampy areas	3.2.4
<i>Casuarina equisetifolia</i> woodland to open forest on foredunes on mainland and islands	3.2.6
<i>Corymbia nesophila</i> ± <i>C. novoguineensis</i> ± <i>Eucalyptus</i> spp. woodland on old stabilised dunes	3.2.8
<i>Eucalyptus phoenicea</i> ± <i>Corymbia nesophila</i> woodland on dunefields around Cape Bedford	3.2.9
Semi-deciduous notophyll vine forest on beach ridges on the east coast	3.2.13
<i>Melaleuca arcana</i> low open forest associated with dune swamps	3.2.14
<i>Melaleuca viridiflora</i> ± <i>Terminalia muelleri</i> low woodland on old beach ridges	3.2.16
<i>Leucopogon yorkensis</i> ± <i>Asteromyrtus angustifolia</i> closed scrub on dunefields	3.2.17
Shrubland to low open heath of <i>Leucopogon</i> spp. on residual beach ridges and sand dunes	3.2.19
<i>Acacia humifusa</i> ± <i>Lithomyrtus obtusa</i> dwarf open heath on dunes and headlands	3.2.22
<i>Neofabricia myrtifolia</i> and <i>Labichea buettneriana</i> dwarf open heath on sand plains	3.2.23
Open herbland of mixed graminoids and forbs on exposed foredunes	3.2.24

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Sparse hermland of mixed herbaceous species on foredunes and beach ridges	3.2.25
Ephemeral and perennial lakes in coastal dunefields	3.2.27
Semi-deciduous notophyll vine forest on beach ridges on coral atolls, shingle cays and sand cays	3.2.28
<i>Pisonia grandis</i> low closed forest restricted to a few scattered sand cays	3.2.29
<i>Pemphis acidula</i> ± <i>Rhizophora stylosa</i> ± <i>Avicennia marina</i> low closed forest on coral atolls, shingle cays and sand cays	3.2.30
<i>Premna serratifolia</i> closed scrub on coral atolls, shingle cays and sand cays	3.2.31
<i>Lepturus repens</i> closed hermland on sand cays	3.2.32
<i>Gahnia sieberiana</i> open to closed heath in drainage swamps in east coast dunefields	3.2.33
Simple evergreen notophyll vine forest ± <i>Wodyetia bifurcata</i> on colluvium of granite ranges	3.3.3
Evergreen mesophyll and/or notophyll vine forest with <i>Archontophoenix</i> spp. on stream banks	3.3.4
Evergreen notophyll vine forest with <i>Melaleuca leucadendra</i> on swamps	3.3.6
Tall semi-deciduous notophyll and/or microphyll vine thicket on colluvial plains	3.3.7
<i>Melaleuca leucadendra</i> ± <i>Eucalyptus tereticornis</i> open forest on alluvium	3.3.11
<i>Melaleuca quinquenervia</i> open forest associated with scattered coastal swamps	3.3.12

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca saligna</i> ± <i>Hakea pedunculata</i> open forest on edges of salt pans	3.3.13
<i>Eucalyptus brassiana</i> woodland on alluvial plains	3.3.15
<i>Corypha utan</i> open woodland on alluvial plains and old beach ridges in Lakefield National Park	3.3.34
Semi-deciduous microphyll vine forest ± <i>Melaleuca</i> spp., associated with sinkholes	3.3.39
<i>Terminalia</i> sp. deciduous vine thicket in depressions in Lakefield area	3.3.40
<i>Vachellia ditricha</i> low open woodland on erosional plains	3.3.44
<i>Eucalyptus chlorophylla</i> ± <i>Melaleuca viridiflora</i> low open woodland on Mitchell River floodplain	3.3.45
<i>Eucalyptus microtheca</i> ± <i>E. chlorophylla</i> low open woodland on Mitchell River alluvia	3.3.46
<i>Melaleuca acacioides</i> ± <i>Hakea pedunculata</i> tall shrubland on marine plains	3.3.51
<i>Asteromyrtus lysicephala</i> ± <i>Jacksonia thesioides</i> open heath on streams on low sandstone plateaus	3.3.54
<i>Imperata cylindrica</i> ± <i>Mnesithea rottboellioides</i> closed-tussock grassland on coastal plains	3.3.57
Grassland and/or sedgeland with <i>Pandanus</i> spp., confined to Torres Strait Islands	3.3.62
Permanent lakes and lagoons, frequently with fringing woodlands or sedgelands	3.3.66
<i>Melaleuca arcana</i> low open forest in swamps	3.3.67

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Semi-deciduous notophyll vine forest and thicket on alluvial plains	3.3.68
<i>Melaleuca dealbata</i> ± <i>Corymbia clarksoniana</i> tall open forest on alluvial plains	3.3.69
<i>Lophostemon suaveolens</i> ± <i>Melaleuca cajuputi</i> subsp. <i>platyphylla</i> ± <i>Pandanus</i> sp. ± <i>Livistona muelleri</i> woodland and open forest on the alluvial plains of the northern Torres Strait Islands	3.3.70
Semi-deciduous notophyll vine forest restricted to lateritic Carnegie Tableland	3.5.3
<i>Corymbia novoguinensis</i> or <i>C. nesophila</i> ± <i>C. tessellaris</i> woodland on sand plains on northern Cape York Peninsula	3.5.5
<i>Melaleuca viridiflora</i> and <i>Asteromyrtus brassii</i> woodland on flat sand plains	3.5.13
<i>Melaleuca stenostachya</i> low open woodland on flat plains	3.5.17
Simple evergreen notophyll vine forest with <i>Eucalyptus pellita</i> on sandstone plateaus	3.5.20
<i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> open forest on coastal ranges and lowlands	3.5.21
<i>Corymbia nesophila</i> ± <i>Eucalyptus crebra</i> ± <i>E. brassiana</i> woodland on plains	3.5.23
<i>Themeda arguens</i> and <i>Dichanthium sericeum</i> closed-tussock grassland on low undulating rises	3.5.30
<i>Corymbia clarksoniana</i> ± <i>Erythrophleum chlorostachys</i> woodland on coastal plains	3.5.31

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Asteromyrtus brassii</i> ± <i>Syzygium angophoroides</i> ± <i>Acmena hemilampra</i> subsp. <i>hemilampra</i> open forest. Residual sand rises and sheets	3.5.32
Semi-deciduous notophyll and/or microphyll vine thicket on isolated lateritic hill slopes	3.7.1
<i>Acacia shirleyi</i> open forest on lateritic knolls	3.7.2
Semi-deciduous notophyll and/or microphyll vine forest on basalt	3.8.2
<i>Eucalyptus leptophleba</i> or <i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> woodland on basalt flows	3.8.3
<i>Imperata cylindrica</i> or <i>Mnesithea rotboellioides</i> tussock grasslands on basalt cones and rises	3.8.4
Semi-deciduous and deciduous notophyll vine forest on the Basaltic Islands of the Torres Strait	3.8.5
<i>Terminalia aridicola</i> subsp. <i>chillagoensis</i> and <i>T.</i> <i>platyphylla</i> open woodland on clay soils	3.9.6
<i>Heteropogon triticeus</i> , <i>Themeda arguens</i> and <i>Sarga</i> <i>plumosum</i> closed-tussock grassland on clay plains	3.9.8
Seepage springs from sandstone or Tertiary plateaus and associated rainforests and vine thickets	3.10.1
Simple evergreen notophyll vine forest with <i>Callitris</i> <i>intratropica</i> on low hills	3.10.3
Deciduous notophyll and/or microphyll vine thicket or forest ± <i>Gyrocarpus americanus</i> or <i>Eucalyptus pellita</i> emergents on sandstone hills and slopes	3.10.5
<i>Eucalyptus similis</i> ± <i>Corymbia nesophila</i> woodland on pediments of sandstone ranges	3.10.8

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Allocasuarina littoralis</i> ± <i>Acacia crassicarpa</i> low woodland on sandstone plateaus	3.10.14
<i>Neofabricia myrtifolia</i> and <i>Acacia calyculata</i> tall open shrubland on sandstone breakaways	3.10.17
Sedgelands, fernlands and closed heathlands associated with springs on sandstone tablelands	3.10.20
Semi-deciduous mesophyll vine forest on coastal ranges	3.11.1
Semi-deciduous mesophyll vine forest on metamorphic ranges in the south	3.11.2
<i>Corymbia nesophila</i> ± <i>Eucalyptus</i> spp. open forest on wetter ranges in south-east	3.11.4
<i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> open forest on metamorphic coastal ranges	3.11.5
<i>Eucalyptus platyphylла</i> ± <i>E. leptophleba</i> ± <i>Corymbia nesophila</i> open forest to woodland on hill slopes	3.11.6
<i>Eucalyptus staigeriana</i> woodland on metamorphic ranges	3.11.14
<i>Eucalyptus tardecidens</i> low woodland on metamorphic plateaus	3.11.16
<i>Themeda triandra</i> tall grassland or <i>Asteromyrtus lysicephala</i> , <i>Neofabricia myrtifolia</i> , <i>Grevillea pteridifolia</i> dwarf open heathlands on headlands and islands	3.11.19
Semi-deciduous mesophyll and/or notophyll vine forest on granite slopes in the McIlwraith Range	3.12.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Araucarian notophyll vine forest on granitic ridges and mountains	3.12.2
Notophyll vine forest of <i>Welchiodendron longivalve</i> and <i>Acacia polystachya</i> on low hills and rises on volcanics	3.12.4
Simple evergreen notophyll vine forest on upper slopes of mountains and ranges	3.12.5
Simple evergreen notophyll vine forest ± <i>Wodyetia bifurcata</i> on colluvium of granite ranges	3.12.6
<i>Eucalyptus brassiana</i> and <i>Corymbia clarksoniana</i> open forest on granite ranges	3.12.7
Evergreen notophyll vine forest dominated by <i>Welchiodendron longivalve</i> on headlands	3.12.20
Deciduous vine thicket ± <i>Wodyetia bifurcata</i> on granite boulders	3.12.22
<i>Acacia brassii</i> low open forest on acid volcanics	3.12.23
<i>Corymbia stockeri</i> ± <i>Eucalyptus crebra</i> low open forest on granite ranges and headlands	3.12.24
<i>Lophostemon suaveolens</i> and <i>Eucalyptus crebra</i> low open forest on granite	3.12.25
<i>Welchiodendron longivalve</i> and <i>Melaleuca viridiflora</i> low woodland on granite ridge crests	3.12.27
<i>Leptospermum purpurascens</i> tall shrubland on acid volcanic hills	3.12.28
<i>Heteropogon triticeus</i> ± <i>Sarga plumosum</i> closed-tussock grassland on continental islands	3.12.29

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Imperata cylindrica</i> ± <i>Mnesithea rottboellioides</i> closed-tussock grassland on steep slopes	3.12.30
<i>Themeda triandra</i> tussock grassland on headlands and islands on acid volcanic rocks	3.12.31
<i>Schizachyrium</i> spp. ± <i>Eriachne</i> spp. tussock grassland on rocky ranges and rock pavements	3.12.32
Granite boulders interspersed with vine thicket	3.12.33
Rock pavements associated with mountains and river beds and some offshore islands	3.12.34
Semi-deciduous mesophyll and/or notophyll vine forest on granite slopes of the Torres Strait subregion	3.12.35
Evergreen to complex evergreen mesophyll to notophyll vine forest and thicket on mountain ranges of Torres Strait Islands	3.12.36
<i>Eucalyptus platyphylla</i> ± <i>Corymbia stockerii</i> ± <i>Corymbia clarksoniana</i> woodland to open woodland on coastal hills	3.12.37
<i>Corymbia clarksoniana</i> ± <i>Corymbia stockerii</i> ± <i>Corymbia nesophila</i> low mixed woodland of Torres Strait Islands	3.12.38

Part 3

Central Queensland Coast Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Sporobolus virginicus</i> tussock grassland on marine sediments	8.1.3
<i>Schoenoplectus subulatus</i> and/or <i>Eleocharis dulcis</i> sedgeland or <i>Paspalum vaginatum</i> tussock grassland	8.1.4
<i>Melaleuca</i> spp. and/or <i>Eucalyptus tereticornis</i> and/or <i>Corymbia tessellaris</i> woodland with a ground stratum of salt tolerant grasses and sedges, usually in a narrow zone adjoining tidal ecosystems	8.1.5
<i>Casuarina equisetifolia</i> woodland and/or sparse hermland to open scrub on foredunes and beaches	8.2.1
Semi-evergreen microphyll vine thicket to vine forest on coastal dunes	8.2.2
<i>Allocasuarina littoralis</i> and/or <i>Leptospermum neglectum</i> and/or <i>Leptospermum polygalifolium</i> and/or <i>Baeckea frutescens</i> shrubland on coastal sand ridges, parabolic dunes and whaleback dunes	8.2.3
Sedgeland, closed heath or <i>Melaleuca</i> spp. open shrubland to open forest on swampy sand plains with peat	8.2.4
Evergreen notophyll <i>Archontophoenix cunninghamiana</i> vine forest in deep depressions and narrow gullies on coastal parabolic dunes	8.2.5
<i>Corymbia tessellaris</i> ± <i>Acacia leptocarpa</i> ± <i>Allocasuarina littoralis</i> ± <i>Banksia integrifolia</i> ± rainforest species open forest on coastal parallel dunes	8.2.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca</i> spp. and/or <i>Lophostemon suaveolens</i> and/or <i>Eucalyptus robusta</i> open forest in wetlands associated with parabolic dunes	8.2.7
Tussock grassland on coastal dunes	8.2.9
Sand blows with bare sand and areas of sparse herbland or shrubland	8.2.10
<i>Melaleuca</i> spp. closed forest in parallel dune swales	8.2.11
<i>Corymbia intermedia</i> and/or <i>Eucalyptus latisinensis</i> and/or <i>Acacia</i> spp. and/or other heath spp. shrublands and woodlands on parallel dunes (subregions 4 and 5)	8.2.12
<i>Banksia integrifolia</i> and/or <i>Corymbia tessellaris</i> and/or <i>Acacia disparrima</i> ± rainforest spp. tall shrubland on Holocene parabolic dunes	8.2.14
Semi-deciduous to evergreen notophyll to mesophyll vine forest ± sclerophyll emergents fringing or in the vicinity of watercourses	8.3.1
<i>Eucalyptus platyphylla</i> and/or <i>Lophostemon suaveolens</i> and/or <i>Corymbia clarksoniana</i> woodland on alluvial plains	8.3.5
<i>Eucalyptus tereticornis</i> and/or <i>Corymbia intermedia</i> (or <i>C. clarksoniana</i>) and/or <i>C. tessellaris</i> ± <i>Lophostemon suaveolens</i> open forest on alluvial levees and lower terraces	8.3.6
<i>Syncarpia glomulifera</i> and/or <i>Eucalyptus portuensis</i> and/or <i>Corymbia intermedia</i> open forest on sandy terrace flats and granite outwash	8.3.8
Semi-deciduous complex notophyll vine forest on perched alluvials in valleys of undulating mountain ranges	8.3.9

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Semi-evergreen to evergreen notophyll vine forest on gently to moderately sloping alluvial fans adjacent to ranges	8.3.10
<i>Eucalyptus tereticornis</i> and/or <i>Corymbia tessellaris</i> and/or <i>Melaleuca</i> spp. woodland on alluvial and marine plains, often adjacent to estuarine areas	8.3.13
<i>Ischaemum australe</i> and/or <i>Imperata cylindrica</i> and/or <i>Sorghum nitidum</i> forma <i>aristatum</i> tussock grassland on drainage channels in gently undulating upland areas	8.3.14
Open water in river channels, waterholes and lagoons, and exposed stream beds and bars	8.3.15
<i>Eucalyptus drepanophylla</i> ± <i>Corymbia clarksoniana</i> ± <i>E. platyphylla</i> ± <i>C. dallachiana</i> ± <i>Melaleuca viridiflora</i> woodland on broad low rises and gently sloping Tertiary sand plains	8.5.3
<i>Eucalyptus exserta</i> and/or <i>Corymbia clarksoniana</i> and/or <i>E. crebra</i> and/or <i>Melaleuca</i> spp. woodland on Tertiary sand plains	8.5.5
<i>Melaleuca viridiflora</i> ± <i>Allocasuarina littoralis</i> woodland on Tertiary sand plains	8.5.6
<i>Melaleuca viridiflora</i> and/or <i>Eucalyptus latisinensis</i> ± <i>Syncarpia glomulifera</i> woodland on Cainozoic sand plains of uncertain age and origin	8.5.7
Evergreen notophyll to complex notophyll vine forest of uplands and highlands on basalt	8.8.1
<i>Eucalyptus latisinensis</i> and/or <i>Corymbia intermedia</i> ± <i>Syncarpia glomulifera</i> woodland on low rises in coastal plains	8.9.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia julifera</i> and/or <i>Eucalyptus</i> spp. ± <i>Corymbia</i> spp. open forest and/or semi-evergreen, simple microphyll low closed forest and/or <i>Heteropogon contortus</i> tussock grassland on slopes of islands on Cretaceous sedimentary rocks	8.10.1
<i>Eucalyptus drepanophylla</i> ± <i>E. platyphylla</i> woodland on hills formed from metamorphosed sediments	8.11.1
Semi-evergreen notophyll to microphyll vine forest of foothills and uplands on metamorphosed sediments	8.11.2
<i>Corymbia tessellaris</i> and/or <i>Eucalyptus tereticornis</i> ± <i>E. drepanophylla</i> open forest on low hills formed from metamorphosed sediments or conglomerate (subregion 2)	8.11.5
<i>Eucalyptus latisinensis</i> and/or <i>E. crebra</i> and/or <i>E. exserta</i> ± <i>Corymbia intermedia</i> ± <i>C. trachyphloia</i> open forest on metamorphosed sediments	8.11.6
<i>Lophostemon confertus</i> and/or <i>Banksia integrifolia</i> and/or <i>Allocasuarina littoralis</i> and/or <i>Xanthorrhoea latifolia</i> shrubland on exposed metamorphic mountain tops	8.11.7
<i>Themeda triandra</i> and/or <i>Heteropogon contortus</i> tussock grassland, or <i>Xanthorrhoea latifolia</i> shrubland with <i>Themeda triandra</i> , on exposed rocky headlands on metamorphosed sediments	8.11.9
<i>Lophostemon</i> spp. and/or <i>Acacia</i> spp. and/or <i>Melaleuca viridiflora</i> and/or <i>Allocasuarina littoralis</i> ± <i>Eucalyptus</i> spp. ± <i>Corymbia</i> spp. tall open shrubland on exposed hill slopes of islands and headlands on metamorphosed sediments	8.11.10

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> and/or <i>E. drepanophylla</i> and/or <i>E. exserta</i> and/or <i>Corymbia clarksoniana</i> and/or <i>C. xanthope</i> and/or <i>Lophostemon confertus</i> low woodland on metamorphics on islands and headlands	8.11.12
<i>Eucalyptus grandis</i> open forest of wet uplands on Mesozoic to Proterozoic igneous rocks (predominantly granite)	8.12.4
<i>Eucalyptus montivaga</i> open forest on plateaus and ridges of high ranges on Mesozoic to Proterozoic igneous rocks	8.12.8
<i>Leptospermum</i> spp. and/or <i>Acacia</i> spp. and/or <i>Lophostemon confertus</i> and/or <i>Allocasuarina littoralis</i> and/or <i>Banksia integrifolia</i> shrubland on plateaus of Cretaceous-Tertiary acid to intermediate volcanics and Mesozoic to Proterozoic igneous rocks	8.12.10
Tussock grassland, or <i>Xanthorrhoea latifolia</i> shrubland, including areas recently colonised by <i>Timonius timon</i> shrubland, on slopes of islands and headlands on Mesozoic to Proterozoic igneous rocks and Tertiary acid to intermediate volcanics	8.12.13
Deciduous to semi-evergreen microphyll vine thicket ± <i>Brachychiton</i> spp. ± <i>Araucaria cunninghamii</i> emergents of foothills and uplands (western areas) on Mesozoic to Proterozoic igneous rocks	8.12.16
Evergreen microphyll to notophyll mossy forest to thicket of ridges and plateaus on highlands to foothills on Mesozoic to Proterozoic igneous rocks	8.12.17
<i>Eucalyptus moluccana</i> woodland on elevated tablelands on Mesozoic to Proterozoic igneous rocks	8.12.23

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tereticornis</i> ± <i>E. tereticornis</i> x <i>E. platyphylla</i> woodland on hillslopes of islands on Mesozoic to Proterozoic igneous rocks	8.12.25
<i>Corymbia tessellaris</i> and/or <i>Eucalyptus tereticornis</i> open forest on hill slopes of islands and near coastal areas on Mesozoic to Proterozoic igneous rocks and Tertiary acid to intermediate volcanics	8.12.26
Semi-evergreen microphyll <i>Acacia fasciculifera</i> , <i>Terminalia</i> spp., <i>Brachychiton</i> spp. vine forest to vine thicket of near-coastal foothills on volcanics (subregion 1)	8.12.28
<i>Allocasuarina littoralis</i> and/or <i>Lophostemon confertus</i> and/or <i>Acacia</i> spp. and/or <i>Grevillea banksii</i> open shrubland on islands and headlands on Mesozoic to Proterozoic igneous and Tertiary acid to intermediate rocks	8.12.29
Evergreen notophyll <i>Ristantia waterhousei</i> mossy forest of uplands on rhyolite	8.12.30

[s 7]

Part 4

Channel Country Bioregion

Column 1**Regional ecosystem****Column 2****Regional
ecosystem
number**

Springs on recent alluvia and fine-grained sedimentary rocks 5.3.23

Acacia calcicola tall shrubland between sand dunes 5.6.3

Acacia peuce low open woodland between dunes 5.7.8

Part 5

Desert Uplands Bioregion

Column 1**Regional ecosystem****Column 2****Regional
ecosystem
number**

Eucalyptus cambageana open woodland on broad stream beds 10.3.5

Acacia excelsa and *Grevillea striata* low open woodland on lake-fringing dunes 10.3.17

Eucalyptus melanophloia open woodland on older lake-fringing dunes 10.3.20

Acacia salicina and *Grevillea striata* low open woodland on sandy alluvial plains 10.3.21

Lysiphyllyum carronii low open woodland on alluvial plains 10.3.26

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia torulosa</i> shrubland or <i>Triodia longiceps</i> hummock grassland on weathered lake dunes	10.3.29
<i>Casuarina cristata</i> woodland on floodplains	10.3.30
Artesian springs emerging on alluvial plains	10.3.31
<i>Acacia harpophylla</i> low woodland on Cainozoic lake beds (subregion 3)	10.4.2
<i>Acacia cambagei</i> woodland on Cainozoic lake beds (subregion 3)	10.4.4
<i>Terminalia oblongata</i> and <i>Lysiphylloum carronii</i> low open woodland on Cainozoic lake beds	10.4.6
<i>Casuarina cristata</i> open woodland on Cainozoic lake beds	10.4.7
<i>Corymbia</i> spp. open woodland on Cainozoic lake beds	10.4.9
<i>Eucalyptus quadricostata</i> open woodland on sandy plateaus	10.5.9
<i>Acacia aneura</i> low open woodland near the margins of sandy plateaus	10.7.6
<i>Eucalyptus exilipes</i> with or without <i>Corymbia leichhardtii</i> open woodland on the perimeter of sandy plateaus	10.7.9
<i>Eucalyptus melanophloia</i> open woodland or <i>Lysiphylloum carronii</i> low open woodland on calcareous sandstones	10.9.5
<i>Melaleuca uncinata</i> dwarf open shrubland on Cretaceous sediments	10.9.7
<i>Archidendropsis basaltica</i> low open woodland on Cretaceous sediments	10.9.8

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus drepanophylla</i> open woodland on sandstone ranges	10.10.3
Springs associated with margins of sandstone plateaus	10.10.6
<i>Eucalyptus cloeziana</i> open woodland on sandstone ranges	10.10.7

Part 6

Einasleigh Uplands Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Permanent or seasonal wetlands frequently fringed by narrow bands of trees and shrubs including <i>Eucalyptus</i> spp. on alluvial plains	9.3.4
<i>Acacia cambagei</i> ± <i>A. harpophylla</i> woodland on cracking clay soils	9.3.9
<i>Eucalyptus chlorophylla</i> ± <i>Corymbia clarksoniana</i> ± <i>Terminalia</i> spp. woodland on alluvial plains	9.3.21
<i>Acacia tephrina</i> open forest on alluvial clay plains	9.3.23
<i>Eucalyptus cambageana</i> woodland with a shrub layer of <i>Eremophila mitchellii</i> , <i>Psydrax oleifolia</i> , <i>Flindersia maculosa</i> and <i>Lysiphyllo</i> spp. on clay lenses in Cainozoic plains	9.4.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus persistens</i> or <i>E. brownii</i> open woodland with a shrub layer of <i>Eremophila mitchellii</i> , <i>Psydrax oleifolia</i> , <i>Flindersia maculosa</i> and <i>Lysiphyllo</i> spp. on clay lenses in Cainozoic plains	9.4.2
<i>Acacia harpophylla</i> and <i>Lysiphyllo carronii</i> open woodland on Cainozoic clays	9.4.3
Semi-evergreen vine thicket on red kandosols on Tertiary plateaus	9.5.2
<i>Melaleuca viridiflora</i> and/or <i>M. stenostachya</i> low open woodland on erosional plains	9.5.14
<i>Melaleuca viridiflora</i> , <i>Grevillea pteridifolia</i> , <i>Allocasuarina littoralis</i> and <i>Callitris intratropica</i> mixed low woodland on Tertiary remnants	9.5.17
<i>Allocasuarina inophloia</i> ± <i>Eucalyptus exserta</i> low open woodland on exposed lateritic surfaces on Tertiary plateaus	9.7.4
<i>Eucalyptus chartaboma</i> ± <i>Corymbia clarksoniana</i> ± <i>Acacia shirleyi</i> woodland on lateritised remnant sand sheets	9.7.6
Semi-evergreen vine thicket on Quaternary basalt soils	9.8.3
Springs associated with basalt and alluvium	9.8.8
<i>Eucalyptus tereticornis</i> and <i>Lophostemon suaveolens</i> woodland ± a shrubby understorey on rocky basalt flows	9.8.10
<i>Excoecaria parvifolia</i> low open woodland on cracking clays on rocky basalt plains	9.8.12
Springs and their associated vegetation on quartzose sandstone, limestone, metamorphic rock and granite	9.10.2

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia trachyphloia</i> and/or <i>Eucalyptus exilipes</i> woodland on remnant sandstone sheets overlying mountain ranges	9.10.4
<i>Eucalyptus similis</i> ± <i>Corymbia erythrophloia</i> open forest on remnant sandstone sheets overlying mountain ranges	9.10.5
<i>Corymbia</i> spp. and <i>Eucalyptus</i> spp. woodland on sandstones of Ngarrabullan	9.10.7
<i>Eucalyptus mediocris</i> and <i>E. cloeziana</i> woodland on sandstones of Ngarrabullan	9.10.8
<i>Acacia johannis</i> low woodland on sandstones of Ngarrabullan	9.10.9
Semi-deciduous vine thicket on metamorphic soils (not limestone)	9.11.9
<i>Eucalyptus cambageana</i> ± <i>E. crebra</i> ± <i>E. brownii</i> woodland on low metamorphic rises	9.11.19
<i>Corymbia setosa</i> ± <i>Eucalyptus crebra</i> low open woodland on metamorphic hills	9.11.21
<i>Eucalyptus melanophloia</i> ± <i>Corymbia erythrophloia</i> ± <i>Terminalia platyptera</i> low woodland on metamorphic hills	9.11.22
<i>Eucalyptus crebra</i> , <i>Corymbia leichhardtii</i> and <i>C. lamprophylla</i> woodland on steep to rugged metamorphic hills	9.11.29
<i>Eucalyptus leptophleba</i> and/or <i>Corymbia terminalis</i> ± <i>C. dallachiana</i> woodland on aprons surrounding limestone outcrops	9.11.32

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Macropteranthes montana</i> low open forest on igneous geologies	9.12.9
<i>Corymbia confertiflora</i> and <i>Eucalyptus crebra</i> ± <i>C. clarksoniana</i> open woodland on rolling igneous hills	9.12.10
<i>Eucalyptus crebra</i> and <i>Corymbia dallachiana</i> ± <i>C. erythrophloia</i> open woodland on pre-Cainozoic basalt loams and flats to undulating plains	9.12.16
<i>Eucalyptus crebra</i> or <i>E. drepanophylla</i> and <i>Corymbia</i> spp. open woodland on flat to undulating country on igneous rocks	9.12.21
<i>Eucalyptus exserta</i> and <i>Lysicarpus angustifolius</i> low open woodland with <i>Triodia bitextura</i> ground layer on sandy soils on igneous rocks	9.12.25
<i>Eucalyptus moluccana</i> ± <i>E. crebra</i> and/or <i>E. granitica</i> woodland on igneous rocks	9.12.26
<i>Eucalyptus similis</i> and <i>E. shirleyi</i> ± <i>E. crebra</i> low open woodland on low granite hills with rocky outcrops	9.12.29
<i>Eucalyptus leptophleba</i> , <i>Corymbia clarksoniana</i> and <i>Eucalyptus crebra</i> ± <i>C. dallachiana</i> woodland on igneous rocks	9.12.31
<i>Melaleuca viridiflora</i> ± <i>Eucalyptus tereticornis</i> ± <i>E. granitica</i> ± <i>Corymbia intermedia</i> low open woodland on igneous uplands	9.12.39
<i>Schizachyrium fragile</i> and <i>Dichanthium sericeum</i> grassland on undulating hills on pre-Cainozoic mafic igneous rocks	9.12.42
Granite and rhyolite boulders and pavements edged with patches of <i>Callitris intratropica</i> ± vine thicket species	9.12.43

[s 7]

Part 7 Gulf Plains Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Beaches and foredunes	2.2.1
<i>Eucalyptus microtheca</i> , <i>Corymbia</i> spp., <i>Lysiphylgium cunninghamii</i> low open woodland with <i>Dichanthium</i> spp. on plains and low rises of texture contrast soils and earths	2.3.8
<i>Excoecaria parvifolia</i> open woodland with sedges in seasonal swamps on grey clay plains	2.3.12
<i>Acacia stenophylla</i> low woodland in seasonal swamps on grey clay plains	2.3.13
<i>Muehlenbeckia florulenta</i> shrubland in channelled depressions in floodplains	2.3.14
<i>Eucalyptus tectifica</i> woodland with <i>Eulalia aurea</i> on plains on solodised solonetz	2.3.19
<i>Eucalyptus microneura</i> woodland in shallow depressions on solodised soils	2.3.35
<i>Eucalyptus platyphylla</i> and <i>Eucalyptus brownii</i> woodland in shallow depressions on plateaus, on podsolics and earths	2.3.37
Sedges in lagoons on plateau surfaces on earths and solodised soils	2.3.38
<i>Callitris glaucophylla</i> woodland on plains on deep sandy soils	2.5.4

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tetrodonta</i> and <i>Corymbia pocillum</i> woodland on earths on low tablelands	2.5.7
Springs associated with quartzose sandstone or lateritised sandstone gullies and gorges	2.10.8

Part 8 Mitchell Grass Downs Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia peuce</i> low open woodland on alluvium	4.3.21
<i>Archidendropsis basaltica</i> , <i>Acacia aneura</i> low open woodland	4.7.3
<i>Acacia cambagei</i> low woodland with scattered shrubs such as <i>Eremophila mitchellii</i> and <i>Geijera parviflora</i> on fresh Cretaceous sediments	4.9.11
<i>Acacia harpophylla</i> tall shrubland with scattered emergent <i>Atalaya hemiglaucha</i> ± <i>Eucalyptus</i> spp. on Cretaceous sediments	4.9.15
<i>Acacia harpophylla</i> ± <i>A. cambagei</i> low woodland on undulating clay plains	4.9.17

[s 7]

Part 9 Mulga Lands Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus populnea, Casuarina cristata or Acacia harpophylla ± Geijera parviflora</i> woodland on clay plains	6.4.3
<i>Eucalyptus populnea, Acacia aneura ± Eremophila mitchellii</i> woodland within <i>A. aneura</i> communities	6.5.3
<i>Eucalyptus populnea ± E. intertexta ± Acacia aneura ± Callitris glaucophylla</i> woodland on Quaternary sediments	6.5.5
<i>Eucalyptus populnea ± E. melanophloia ± Callitris glaucophylla ± Acacia aneura</i> woodland on sand plains	6.5.17
Springs associated with lateritised sandstone	6.7.18
Scattered <i>Acacia aneura</i> around granite boulders	6.12.1

Part 10 New England Tableland Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> fringing open forest	13.3.5
Sedgelands on alluvial deposits in igneous landscapes	13.3.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus laevopinea</i> open forest on metamorphics	13.11.2
<i>Eucalyptus crebra</i> woodland on metamorphics	13.11.3
<i>Eucalyptus sideroxylon</i> , <i>E. fibrosa</i> subsp. <i>nubila</i> open forest on metamorphics	13.11.5
Low microphyll vine forest on metamorphics	13.11.7
<i>Eucalyptus melliodora</i> and/or <i>Eucalyptus microcarpa</i> / <i>E. moluccana</i> woodland on metamorphics	13.11.8
<i>Eucalyptus scoparia</i> woodland on igneous rocks	13.12.3
<i>Eucalyptus caliginosa</i> , <i>E. tereticornis</i> open forest on igneous rocks	13.12.4
Shrubland on igneous rocks	13.12.6

Part 11 Northwest Highlands Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia georginae</i> low woodland and/or low open woodland on clay plains	1.3.3
Perennial watercourses and associated alluvium	1.3.9
<i>Eucalyptus miniata</i> woodland on red earths on laterised plateaus	1.5.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Mixed eucalypt woodland on sandy plains	1.5.2
Mixed shrubby woodland on low rocky hills on Tertiary limestones	1.9.2
<i>Eucalyptus leucophylla</i> low open woodland on hillocks on Mesozoic claystones	1.9.3
Spring wetlands on undeformed fine-grained sedimentary rock	1.9.8
Springs mostly associated with quartzose sandstone and fine-grained sedimentary rocks (limestone)	1.10.6
Springs associated with metamorphic rocks	1.11.5

Part 12 South East Queensland Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Casuarina glauca</i> woodland on margins of marine clay plains	12.1.1
Notophyll vine forest on parabolic high dunes	12.2.1
Microphyll to notophyll vine forest on beach ridges	12.2.2
Araucarian vine forest on parabolic high dunes	12.2.3
<i>Syncarpia hillii</i> , <i>Lophostemon confertus</i> tall open to closed forest on parabolic high dunes	12.2.4

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Open or dry heath on dunes and beaches	12.2.13
Sand blows largely devoid of vegetation	12.2.16
Mixed closed-tussock grassland to closed hermland on coral, shingle and sand cays	12.2.17
<i>Abutilon albescens</i> ± <i>Wollastonia biflora</i> low shrubland, restricted to coral, shingle and sand cays	12.2.18
<i>Argusia argentea</i> low woodland, restricted to coral, shingle and sand cays	12.2.19
<i>Pandanus tectorius</i> open woodland ± <i>Celtis paniculata</i> and <i>Pisonia grandis</i> , restricted to established cays	12.2.20
<i>Pisonia grandis</i> low closed forest, restricted to established cays	12.2.21
<i>Eucalyptus grandis</i> tall open forest on alluvial plains	12.3.2
<i>Melaleuca quinquenervia</i> , <i>Eucalyptus robusta</i> woodland on coastal alluvium	12.3.4
Swamps with <i>Cyperus</i> spp., <i>Schoenoplectus</i> spp. and <i>Eleocharis</i> spp.	12.3.8
<i>Eucalyptus nobilis</i> open forest on alluvial plains	12.3.9
<i>Eucalyptus tereticornis</i> ± <i>Eucalyptus siderophloia</i> , <i>Corymbia intermedia</i> open forest on alluvial plains, usually near coast	12.3.11
<i>Banksia aemula</i> low woodland on alluvial plains, usually near coast	12.3.14
<i>Corymbia intermedia</i> , <i>Syncarpia glomulifera</i> open forest on granite outwash	12.3.15

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus portuensis</i> , <i>Corymbia intermedia</i> open forest on remnant Tertiary surfaces, usually deep red soils	12.5.5
<i>Eucalyptus hallii</i> open woodland on complex of remnant Tertiary surface and Tertiary sedimentary rocks	12.5.8
Sedgeland to heathland in low lying areas on complex of remnant Tertiary surface and Tertiary sedimentary rocks	12.5.9
<i>Eucalyptus racemosa</i> , <i>E. latisinensis</i> ± <i>Corymbia gummifera</i> , <i>C. intermedia</i> , <i>E. bancroftii</i> woodland with heathy understorey on remnant Tertiary surfaces	12.5.12
<i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> and/or <i>Corymbia trachyphloia</i> woodland on jump-ups	12.7.1
<i>Eucalyptus rhombica</i> , <i>Corymbia trachyphloia</i> woodland on jump-ups	12.7.2
<i>Eucalyptus oreades</i> tall open forest on Cainozoic igneous rocks	12.8.2
Simple microphyll fern forest with <i>Nothofagus moorei</i> on Cainozoic igneous rocks	12.8.6
Simple microphyll fern thicket with <i>Acmena smithii</i> on Cainozoic igneous rocks	12.8.7
<i>Eucalyptus saligna</i> or <i>E. grandis</i> tall open forest on Cainozoic igneous rocks	12.8.8
<i>Eucalyptus laevopinea</i> tall open forest on Cainozoic igneous rocks	12.8.10
<i>Eucalyptus dunnii</i> tall open forest on Cainozoic igneous rocks	12.8.11

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus obliqua</i> tall open forest on Cainozoic igneous rocks	12.8.12
Araucarian complex microphyll vine forest on Cainozoic igneous rocks	12.8.13
<i>Poa labillardieri</i> var. <i>labillardierei</i> grassland on Cainozoic igneous rocks	12.8.15
<i>Eucalyptus crebra</i> ± <i>E. melliodora</i> , <i>E. tereticornis</i> woodland on Cainozoic igneous rocks	12.8.16
Simple notophyll vine forest with <i>Ceratopetalum apetalum</i> on Cainozoic igneous rocks	12.8.18
Heath and rock pavement with scattered shrubs or open woodland on Cainozoic igneous hills and mountains	12.8.19
Shrubby woodland with <i>Eucalyptus racemosa</i> or <i>E. dura</i> on Cainozoic igneous rocks	12.8.20
Open forest with <i>Eucalyptus acmenoides</i> or <i>E. helidonica</i> on Cainozoic igneous rocks especially trachyte	12.8.25
<i>Corymbia trachyphloia</i> and <i>Eucalyptus major</i> woodland on igneous rocks	12.8.26
Tall open forest often with <i>Eucalyptus resinifera</i> , <i>E. grandis</i> , <i>E. robusta</i> and <i>Corymbia intermedia</i> on sedimentary rocks. Coastal	12.9–10.1
<i>Eucalyptus moluccana</i> open forest on sedimentary rocks	12.9–10.3
<i>Eucalyptus crebra</i> ± <i>E. tereticornis</i> , <i>Corymbia tessellaris</i> , <i>Angophora</i> spp. and <i>E. melanophloia</i> woodland on sedimentary rocks	12.9–10.7

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Shrubland and low woodland on sandstone lithosols	12.9–10.9
<i>Melaleuca nodosa</i> low open forest on sedimentary rocks	12.9–10.10
<i>Eucalyptus corynoides</i> woodland on sedimentary rocks	12.9–10.13
Araucarian microphyll to notophyll vine forest on Cainozoic and Mesozoic sediments	12.9–10.16
<i>Angophora leiocarpa</i> , <i>Eucalyptus crebra</i> woodland on sedimentary rocks	12.9–10.18
<i>Eucalyptus montivaga</i> woodland on sedimentary rocks	12.9–10.20
Closed sedgeland and/or shrubland on sedimentary rocks. Coastal parts	12.9–10.22
<i>Eucalyptus melanoleuca</i> open forest on sedimentary rocks	12.9–10.23
<i>Eucalyptus suffulgens</i> open forest on sedimentary rocks	12.9–10.24
Semi-evergreen vine thicket on metamorphics ± interbedded volcanics	12.11.4
<i>Eucalyptus melanophloia</i> , <i>E. crebra</i> woodland on metamorphics ± interbedded volcanics	12.11.8
<i>Eucalyptus tereticornis</i> open forest on metamorphics ± interbedded volcanics, usually higher altitudes	12.11.9
Araucarian complex microphyll vine forest on metamorphics ± interbedded volcanics, usually in northern half of bioregion	12.11.12
Semi-evergreen vine thicket on metamorphics ± interbedded volcanics, usually in northern half of bioregion	12.11.13

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> , <i>E. tereticornis</i> , <i>Corymbia intermedia</i> woodland on metamorphics ± interbedded volcanics	12.11.14
<i>Eucalyptus tereticornis</i> , <i>Corymbia intermedia</i> open woodland with <i>Xanthorrhoea johnsonii</i> understorey on serpentinite	12.11.15
<i>Eucalyptus acmenoides</i> or <i>E. portuensis</i> open forest on metamorphics ± interbedded volcanics	12.11.17
<i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> woodland on metamorphics ± interbedded volcanics	12.11.19
<i>Corymbia intermedia</i> and <i>Lophostemon suaveolens</i> woodland on metamorphics ± interbedded volcanics	12.11.20
<i>Allocasuarina luehmannii</i> and <i>Melaleuca nervosa</i> woodland on metamorphics ± interbedded volcanics	12.11.21
Simple notophyll vine forest usually with abundant <i>Archontophoenix cunninghamiana</i> (gully vine forest) on Mesozoic to Proterozoic igneous rocks	12.12.1
<i>Eucalyptus montivaga</i> open forest on Mesozoic to Proterozoic igneous rocks	12.12.6
<i>Eucalyptus melanophloia</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.8
<i>Eucalyptus dura</i> woodland usually on rocky peaks on Mesozoic to Proterozoic igneous rocks	12.12.9
Shrubland of rocky peaks on Mesozoic to Proterozoic igneous rocks	12.12.10
<i>Eucalyptus tereticornis</i> , <i>Corymbia intermedia</i> , <i>E. crebra</i> ± <i>Lophostemon suaveolens</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.12

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus racemosa</i> subsp. <i>racemosa</i> ± <i>Lophostemon confertus</i> , <i>Syncarpia glomulifera</i> , <i>Eucalyptus acmenoides</i> woodland usually on rocky near-coastal areas on Mesozoic to Proterozoic igneous rocks	12.12.14
Semi-evergreen vine thicket on Mesozoic to Proterozoic igneous rocks, north of bioregion	12.12.18
Vegetation complex of rocky headlands on Mesozoic to Proterozoic igneous rocks	12.12.19
<i>Eucalyptus saligna</i> tall open forest on Mesozoic to Proterozoic igneous rocks	12.12.20
<i>Corymbia intermedia</i> , <i>E. exserta</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.21
<i>Eucalyptus decolor</i> , <i>E. portuensis</i> or <i>E. acmenoides</i> open forest on Mesozoic to Proterozoic igneous rocks	12.12.22
<i>Corymbia trachyphloia</i> , <i>Eucalyptus crebra</i> and <i>Callitris endlicheri</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.27
<i>Eucalyptus moluccana</i> open forest on Mesozoic to Proterozoic igneous rocks	12.12.28

Part 13 Wet Tropics Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Sporobolus virginicus</i> grassland, samphire open formland to sparse formland and bare saltfans on plains adjacent to mangroves	7.1.2
<i>Schoenoplectus subulatus</i> and/or <i>Eleocharis dulcis</i> sparse sedgeland, or <i>Melaleuca quinquenervia</i> low open forest, in swamps which fluctuate periodically between freshwater and estuarine	7.1.3
Mangrove and vine forest closed forest of the brackish zone	7.1.4
<i>Melaleuca viridiflora</i> or <i>Melaleuca</i> spp. ± <i>Acacia</i> spp. ± mangrove spp. woodland on plains adjacent to mangroves	7.1.5
Notophyll to microphyll vine forest on sands of beach origin	7.2.2
<i>Corymbia tessellaris</i> and/or <i>Acacia crassicarpa</i> and/or <i>C. intermedia</i> and/or <i>C. clarksoniana</i> woodland to closed forest on beach ridges (predominantly Holocene)	7.2.3
<i>Eucalyptus</i> spp. (often <i>E. pellita</i> or <i>Corymbia intermedia</i>) open forest and/or <i>Lophostemon suaveolens</i> open forest on swampy sand plains and Pleistocene beach ridges	7.2.4
Mesophyll to notophyll vine forest of <i>Syzgium forte</i> subsp. <i>forte</i> on sands of beach origin	7.2.5
Mosaic of clumps of notophyll vine forest, sclerophyll spp. shrubland and open woodland, and bare sand blows on aeolian dunes	7.2.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Casuarina equisetifolia</i> ± <i>Corymbia tessellaris</i> open forest ± groved vine forest shrublands on strand and foredunes	7.2.7
<i>Melaleuca leucadendra</i> open forest to woodland on sands of beach origin	7.2.8
<i>Melaleuca quinquenervia</i> shrubland to closed forest, or <i>Lepironia articulata</i> open to closed sedgeland, on dune swales and swampy sand plains of beach origin	7.2.9
Shrubland, sedgeland and heath complex with <i>Thryptomene oligandra</i> and/or <i>Asteromyrtus</i> spp. ± <i>Melaleuca quinquenervia</i> on sand plains of beach origin	7.2.10
<i>Melaleuca viridiflora</i> ± <i>Lophostemon suaveolens</i> ± emergent <i>Eucalyptus</i> spp. woodland to open forest, or <i>Melaleuca viridiflora</i> var. <i>attenuata</i> open forest to woodland, on swampy sand plains of beach origin	7.2.11
Grasslands and sedgelands ± <i>Melaleuca</i> spp. within volcanic craters, often on peat	7.3.2
Mesophyll vine forest with <i>Archontophoenix alexandrae</i> on poorly drained alluvial plains	7.3.3
Mesophyll vine forest with <i>Licuala ramsayi</i> on poorly drained alluvial plains and alluvial areas of uplands	7.3.4
Simple–complex mesophyll to notophyll vine forest on moderate to poorly drained alluvial plains of moderate fertility	7.3.10
<i>Corymbia nesophila</i> open forest to woodland on alluvium	7.3.13

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus leptophleba</i> ± <i>Corymbia clarksoniana</i> ± <i>Melaleuca dealbata</i> woodland to open forest on alluvium in low rainfall areas of the west and north	7.3.14
<i>Corymbia intermedia</i> or <i>C. tessellaris</i> ± <i>Eucalyptus tereticornis</i> open forest (or vine forest with these species as emergents) on well-drained alluvium	7.3.19
<i>Corymbia intermedia</i> and <i>Syncarpia glomulifera</i> , or <i>C. intermedia</i> and <i>Eucalyptus pellita</i> , or <i>S. glomulifera</i> and <i>Allocasuarina</i> spp., or <i>E. cloeziana</i> , or <i>C. torelliana</i> open forest (or vine forest with these emergents) on alluvial fans at the base of ranges	7.3.20
<i>Eucalyptus portuensis</i> ± <i>Corymbia intermedia</i> open forest to woodland on alluvium on alluvial fans at the base of ranges	7.3.21
<i>Melaleuca leucadendra</i> ± vine forest species open forest to closed forest on alluvium fringing streams	7.3.25
<i>Casuarina cunninghamiana</i> woodland to open forest on alluvium fringing streams	7.3.26
Rivers and streams including riparian herbfield and shrubland on river and stream bed alluvium and rock within stream beds	7.3.28
Sedgelands and grasslands of permanently and semi-permanently inundated swamps, including areas of open water	7.3.29
<i>Lepironia articulata</i> sedgeland to open sedgeland of permanently to semi-permanently inundated peat swamps of alluvial plains	7.3.31
Open water and narrow shoreline sedge fringes of lakes in volcanic craters	7.3.33

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Complex mesophyll vine forest or simple notophyll vine forest of high rainfall, cloudy uplands on alluvium	7.3.36
Complex notophyll vine forest with emergent <i>Agathis robusta</i> on alluvial fans	7.3.38
<i>Eucalyptus tereticornis</i> ± <i>E. platyphylla</i> ± <i>Corymbia intermedia</i> ± <i>Lophostemon suaveolens</i> open woodland to open forest and associated sedgelands and grasslands on broad drainage depressions of uplands	7.3.39
<i>Eucalyptus grandis</i> open forest to woodland (or vine forest with emergent <i>E. grandis</i>) on alluvium	7.3.42
<i>Eucalyptus tereticornis</i> open forest to woodland on uplands on well-drained alluvium	7.3.43
<i>Allocasuarina littoralis</i> , <i>Corymbia intermedia</i> and <i>Lophostemon suaveolens</i> open forest on poorly drained alluvium	7.3.47
<i>Eucalyptus portuensis</i> and <i>E. drepanophylla</i> ± <i>Corymbia intermedia</i> ± <i>C. citriodora</i> open woodland to open forest on dry uplands on alluvium	7.3.48
Notophyll vine forest on rubble terraces of streams	7.3.49
<i>Melaleuca fluviatilis</i> ± vine forest species open forest to closed forest on alluvium fringing streams	7.3.50
<i>Eucalyptus tereticornis</i> , <i>Corymbia intermedia</i> and <i>E. reducta</i> woodland to open forest of uplands on weathered soils of a remnant surface	7.5.1
<i>Eucalyptus portuensis</i> ± <i>Corymbia intermedia</i> open forest to woodland of uplands on weathered soils of a remnant surface	7.5.2

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus portuensis</i> , <i>Corymbia citriodora</i> and <i>E. drepanophylla</i> woodland to open forest of uplands on weathered soils of a remnant surface	7.5.3
<i>Corymbia intermedia</i> or <i>Melaleuca viridiflora</i> woodland to open forest of uplands on weathered soils of a remnant surface	7.5.4
<i>Eucalyptus tereticornis</i> open forest to tall open forest and associated grasslands, predominantly on basalt uplands	7.8.7
<i>Eucalyptus tereticornis</i> , <i>E. reducta</i> ± <i>Angophora floribunda</i> open forest to woodland on basalt	7.8.8
<i>Eucalyptus tereticornis</i> , <i>E. drepanophylla</i> (or <i>E. granitica</i>), <i>E. portuensis</i> , <i>Corymbia intermedia</i> woodland to open forest, or <i>E. moluccana</i> woodland to open forest, of uplands and highlands on basalt	7.8.10
Closed vineland of wind-disturbed vine forest on basalt foothills and coastal ranges	7.8.11
Complex notophyll vine forest dominated by <i>Backhousia bancroftii</i> on basaltic terraces and scree slopes of the North Johnstone River	7.8.12
Simple notophyll vine forest of <i>Blepharocarya involucrigera</i> of high rainfall, cloudy uplands on basalt	7.8.13
Complex notophyll vine forest with emergent <i>Agathis robusta</i> , on basalt	7.8.14
<i>Eucalyptus grandis</i> open forest to woodland (or vine forest with <i>E. grandis</i> emergents) on basalt	7.8.15
<i>Eucalyptus resinifera</i> open forest to woodland on basalt	7.8.16

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus portuensis</i> and <i>Corymbia intermedia</i> ± <i>C. citriodora</i> open forest to woodland on basalt	7.8.17
<i>Corymbia intermedia</i> and/or <i>Lophostemon suaveolens</i> ± <i>Allocasuarina torulosa</i> open forest to woodland on basalt	7.8.18
Notophyll or mesophyll vine forest with <i>Archontophoenix alexandrae</i> or <i>Licuala ramsayi</i> on metamorphics	7.11.2
Semi-deciduous mesophyll vine forest on moist and dry metamorphic foothills	7.11.3
<i>Syncarpia glomulifera</i> ± <i>Eucalyptus pellita</i> open forest of deep soils on metamorphics	7.11.6
<i>Acacia polystachya</i> woodland to closed forest, or <i>Acacia mangium</i> and <i>Acacia celsa</i> open forest to closed forest, on metamorphics	7.11.8
<i>Acacia celsa</i> open forest to closed forest on metamorphics	7.11.10
<i>Corymbia torelliana</i> open forest, usually with a vine forest element, on metamorphics	7.11.13
<i>Eucalyptus grandis</i> open forest to woodland, or <i>Corymbia intermedia</i> , <i>E. pellita</i> and <i>E. grandis</i> open forest to woodland (or vine forest with these species as emergents), on metamorphics	7.11.14
<i>Eucalyptus portuensis</i> and <i>Corymbia intermedia</i> open forest to woodland on metamorphics of foothills and uplands	7.11.16

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia intermedia</i> and/or <i>C. tessellaris</i> ± <i>Eucalyptus tereticornis</i> open forest to woodland (or vine forest with these species as emergents) on coastal metamorphic headlands and foothills	7.11.18
<i>Corymbia intermedia</i> and/or <i>Lophostemon suaveolens</i> open forest to woodland on uplands on metamorphics	7.11.19
Complex mesophyll vine forest on fertile, well-drained metamorphics of very wet and wet footslopes	7.11.23
Closed vineland of wind-disturbed vine forest of metamorphic slopes, often steep and exposed	7.11.24
Simple–complex mesophyll to notophyll vine forest on amphibolites of the very wet lowlands and foothills	7.11.25
<i>Allocasuarina littoralis</i> and <i>Syncarpia glomulifera</i> open shrubland to closed scrub or <i>Bombax ceiba</i> and <i>Cochlospermum gillivraei</i> open woodland or <i>Acacia</i> spp. shrubland on metamorphic rock pavements	7.11.26
Simple microphyll vine-fern forest or microphyll vine-sedge forest of wet metamorphic uplands and highlands	7.11.27
Wind-sheared notophyll vine forest of exposed metamorphic ridge crests and steep slopes	7.11.28
Microphyll to notophyll vine forests with <i>Ceratopetalum virchowii</i> and/or <i>Uromyrtus</i> <i>metrosideros</i> , <i>Flindersia bourjotiana</i> , <i>F. pimenteliana</i> and <i>Beilschmiedia oligandra</i> of moist uplands on sharply undulating metamorphics	7.11.29
Simple notophyll vine forest of <i>Blepharocarya</i> <i>involucrigera</i> on metamorphics	7.11.30

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus resinifera</i> ± <i>Eucalyptus portuensis</i> ± <i>Syncarpia glomulifera</i> open forest to woodland (or vine forest with these species as emergents) on metamorphics	7.11.31
<i>Syncarpia glomulifera</i> and/or <i>Allocasuarina</i> spp. ± heathy understorey, woodland to tall woodland to open forest (or vine forest with these species as emergents) on steep rocky metamorphic slopes with shallow soils	7.11.32
<i>Eucalyptus reducta</i> open forest to woodland on metamorphics	7.11.33
Complex of shrublands, low heathy or shrubby woodland and low open forest, with <i>Corymbia</i> <i>tessellaris</i> and <i>C. intermedia</i> or <i>Melaleuca viridiflora</i> , <i>Allocasuarina</i> spp. and <i>Acacia</i> spp. on metamorphic coastal headlands and islands	7.11.34
<i>Allocasuarina littoralis</i> , <i>Corymbia intermedia</i> , <i>Lophostemon suaveolens</i> , <i>Xanthorrhoea johnsonii</i> shrubland on serpentinite foothills with deep red soils	7.11.36
<i>Eucalyptus drepanophylla</i> and <i>Corymbia clarksoniana</i> or <i>C. erythrophloia</i> woodland to open forest on dry uplands on metamorphics between Tolga and Mount Molloy	7.11.37
<i>Lophostemon confertus</i> low woodland to low closed forest ± <i>Acacia celsa</i> , <i>Syncarpia glomulifera</i> and <i>Allocasuarina</i> spp. on steep metamorphic slopes	7.11.38
<i>Themeda triandra</i> , or <i>Imperata cylindrica</i> , <i>Sorghum</i> <i>nitidum</i> and <i>Mnesithea rottboellioides</i> closed-tussock grassland of metamorphic headlands and near-coastal hills	7.11.39

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Complex of sclerophyll communities dominated by <i>Syncarpia glomulifera</i> or <i>Melaleuca</i> spp., or sedges, ferns or microphyll vine forest with <i>Trochocarpa bellendenkerensis</i> on highlands, on quartzite or associated metamorphics	7.11.40
<i>Melaleuca viridiflora</i> , <i>M. monantha</i> , <i>Acacia flavescens</i> and <i>Grevillea</i> spp. shrubland with emergent <i>Corymbia clarksoniana</i> , or open woodland of <i>Eucalyptus drepanophylla</i> with <i>M. monantha</i> or <i>Callitris intratropica</i> , on metamorphics	7.11.41
<i>Eucalyptus tereticornis</i> , <i>Pandanus</i> sp., <i>Lophostemon suaveolens</i> , <i>Melaleuca dealbata</i> and <i>E. pellita</i> woodland to open forest of perched drainage areas on metamorphics	7.11.42
<i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> open forest to woodland on metamorphic coastal lowlands and foothills	7.11.43
<i>Eucalyptus tereticornis</i> open forest to woodland on coastal metamorphic foothills	7.11.44
<i>Eucalyptus cloeziana</i> open forest on metamorphics	7.11.45
<i>Eucalyptus portuensis</i> open forest, often with <i>Corymbia nesophila</i> , on near-coastal metamorphic foothills north of the Daintree River	7.11.46
<i>Corymbia nesophila</i> open forest on moderate to steep metamorphic slopes	7.11.47
<i>Melaleuca viridiflora</i> ± <i>Corymbia clarksoniana</i> ± <i>Eucalyptus platyphylла</i> woodland to open forest on metamorphics	7.11.48

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus leptophleba</i> , <i>Corymbia clarksoniana</i> and <i>E. platyphylla</i> open forest to woodland on metamorphic foothills	7.11.49
<i>Eucalyptus platyphylla</i> ± <i>E. drepanophylla</i> ± <i>Corymbia</i> spp. open woodland to open forest on metamorphics	7.11.50
Notophyll or mesophyll vine forest with <i>Archontophoenix alexandrae</i> or <i>Licuala ramsayi</i> , on granites and rhyolites	7.12.2
<i>Syncarpia glomulifera</i> ± <i>Eucalyptus pellita</i> open forest of granites and rhyolites on deep soils	7.12.4
<i>Eucalyptus pellita</i> ± <i>Corymbia intermedia</i> open forest, or <i>Acacia mangium</i> and <i>Lophostemon suaveolens</i> open forest (or vine forest with these species as emergents), on granites and rhyolites	7.12.5
Semi-deciduous mesophyll vine forest on granites and rhyolites of the moist and dry lowlands and foothills	7.12.6
<i>Acacia celsa</i> open forest to closed forest on granites and rhyolites	7.12.9
Notophyll vine forest with emergent <i>Araucaria cunninghamii</i> on moist and dry granite foothills and uplands	7.12.10
<i>Acacia mangium</i> and <i>A. celsa</i> open forest to closed forest or <i>A. polystachya</i> woodland to closed forest of granite and rhyolite foothills	7.12.12
<i>Acacia melanoxylon</i> and <i>A. celsa</i> closed forest of cloudy wet uplands and highlands on granites and rhyolites	7.12.13

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia torelliana</i> open forest usually with a well developed simple notophyll vine forest element on granites and rhyolites	7.12.17
Simple microphyll vine-fern thicket of windswept exposed peaks on granites	7.12.20
<i>Corymbia intermedia</i> and/or <i>C. tessellaris</i> ± <i>Eucalyptus tereticornis</i> open forest to tall open forest to woodland (or vine forest with these species as emergents) on coastal granite and rhyolite headlands and near-coastal foothills	7.12.23
<i>Eucalyptus cloeziana</i> open forest to woodland on granite and rhyolite, often on poorly drained soils	7.12.25
<i>Corymbia nesophila</i> woodland to open forest on granites	7.12.33
<i>Eucalyptus portuensis</i> , <i>E. tereticornis</i> , <i>Corymbia intermedia</i> woodland on granites and rhyolites in the Kirrama-Oak Hills area	7.12.35
Rock pavements and seepage areas of wet lowlands, uplands and highlands of the eastern escarpment and central range (excluding Hinchinbrook Island and Bishops Peak) on granite and rhyolite with <i>Allocasuarina</i> spp. shrublands and/or sedgelands	7.12.37
Deciduous microphyll vine forest and/or blue-green algae-covered granite and rhyolite boulderfields	7.12.38
Complex mesophyll vine forest on fertile, well-drained granites and rhyolites of very wet and wet lowlands, foothills and uplands	7.12.39
Closed vineland of wind-disturbed vine forest on granites and rhyolites	7.12.40

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Podocarpus grayae</i> , <i>Callitris endlicheri</i> and <i>Acacia celsa</i> heathland/shrubland on steep rocky granite slopes of the Hinchinbrook Island uplands and highlands	7.12.41
Notophyll vine forest with <i>Flindersia brayleyana</i> and <i>Argyrodendron polyandrum</i> on granite uplands of Great Palm Island	7.12.42
Simple notophyll vine forest dominated by <i>Stockwellia quadrifida</i> on granites	7.12.43
Simple notophyll vine forest dominated by <i>Blepharocarya involucrigera</i> on granites	7.12.44
Simple notophyll vine forest dominated by <i>Dryadodaphne trachyphloia</i> on granites	7.12.45
Microphyll vine forest with <i>Gossia bidwillii</i> ± <i>Araucaria cunninghamii</i> on steep granite talus and boulder slopes of the Palm Islands	7.12.46
Notophyll-microphyll semi-evergreen vine forest with <i>Argyrodendron polyandrum</i> emergents on rhyolites	7.12.47
Wind-sheared notophyll vine forest on exposed granite and rhyolite ridge crests and steep slopes	7.12.48
Notophyll vine forest and thicket with <i>Planchonella euphlebia</i> and <i>Podocarpus grayae</i> on granites	7.12.49
Simple microphyll vine-fern forest of highlands on granites and rhyolites	7.12.50
<i>Eucalyptus resinifera</i> , <i>Syncarpia glomulifera</i> , <i>E. portuensis</i> , <i>Corymbia abergiana</i> ± <i>C. leptoloma</i> woodland of rocky hills on granite and rhyolite in the Paluma-Seaview (south-west) subregion	7.12.51

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus resinifera, Corymbia intermedia, Allocasuarina littoralis, Syncarpia glomulifera, E. drepanophylla ± E. reducta</i> woodland on granite and rhyolite in the dry to moist rainfall zone	7.12.52
Complex of shrubland and low open forest on wind-exposed granite and rhyolite coastal headlands and islands on skeletal soils	7.12.54
<i>Eucalyptus leptophleba</i> woodland to open forest of dry foothills and uplands on granites and rhyolites	7.12.55
<i>Corymbia tessellaris, C. clarksoniana</i> grassy woodland, open woodland and grassland on shallow soils on granites on the Palm Islands	7.12.56
Shrubland and low woodland mosaic with <i>Syncarpia glomulifera, Corymbia abergiana, Eucalyptus portuensis, Allocasuarina littoralis</i> and <i>Xanthorrhoea johnsonii</i> on uplands and highlands on granites	7.12.57
<i>Eucalyptus reducta ± E. granitica ± Corymbia dimorpha ± C. citriodora</i> woodland to open forest on granites and rhyolites	7.12.58
<i>Eucalyptus leptophleba</i> and <i>Corymbia clarksoniana</i> open forest to woodland on foothills on granites and rhyolites	7.12.59
<i>Melaleuca viridiflora ± Corymbia clarksoniana ± Eucalyptus platyphyllo</i> woodland to open forest on granites and rhyolites	7.12.60
<i>Eucalyptus</i> spp. and/or <i>Corymbia stockeri ± C. hylandii ± Syncarpia glomulifera ± E. portuensis</i> woodland on dry granite hill slopes in the north-west of the bioregion	7.12.62

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus moluccana</i> woodland on granites and rhyolites	7.12.63
<i>Xanthorrhoea</i> spp., <i>Allocasuarina littoralis</i> , <i>Banksia plagiocarpa</i> ± <i>Leptospermum polygalifolium</i> ± <i>Rhodomyrtus trineura</i> subsp. <i>trineura</i> heathland and associated rock pavements of granite uplands and highlands of Hinchinbrook Island and near Bishops Peak	7.12.64
<i>Lophostemon confertus</i> low shrubland or low closed forest on exposed rocky slopes on granites and rhyolites	7.12.66
<i>Gleichenia dicarpa</i> , <i>Gahnia sieberiana</i> , <i>Lycopodiella cernua</i> and <i>Lycopodium deuterodensum</i> closed fernland of granite highlands on Thornton Peak and Mt Bartle Frere	7.12.67
Complex notophyll vine forest of cloudy moist to wet highlands on granites	7.12.68
<i>Eucalyptus drepanophylla</i> and/or <i>E. granitica</i> ± <i>Corymbia clarksoniana</i> ± <i>C. erythrophloia</i> woodland on uplands on granites and rhyolites	7.12.69

Schedule 3 Least concern regional ecosystems

section 8(3) and (6)

Part 1 Brigalow Belt Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Sporobolus virginicus</i> grassland on marine clay plains	11.1.1
Samphire forbland on marine clay plains	11.1.2
Mangrove forest and/or woodland on marine clay plains	11.1.4
<i>Corymbia-Melaleuca</i> woodland complex of beach ridges and swales	11.2.5
<i>Acacia cambagei</i> woodland on alluvial plains	11.3.5
<i>Eucalyptus melanophloia</i> woodland on alluvial plains	11.3.6
<i>Corymbia</i> spp. woodland on alluvial plains	11.3.7
<i>Acacia argyrodendron</i> woodland on alluvial plains	11.3.8
<i>Eucalyptus platyphylla</i> , <i>Corymbia</i> spp. woodland on alluvial plains	11.3.9
<i>Eucalyptus brownii</i> woodland on alluvial plains	11.3.10
<i>Melaleuca viridiflora</i> , <i>M. argentea</i> ± <i>M. dealbata</i> woodland on alluvial plains	11.3.12

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus</i> spp., <i>Angophora</i> spp., <i>Callitris</i> spp. woodland on alluvial plains	11.3.14
<i>Eucalyptus largiflorens</i> ± <i>Acacia cambagei</i> ± <i>A. harpophylla</i> woodland to low open woodland on alluvial plains	11.3.16
<i>Eucalyptus populnea</i> , <i>Callitris glaucophylla</i> , <i>Allocasuarina luehmannii</i> shrubby woodland on alluvium	11.3.18
<i>Callitris glaucophylla</i> , <i>Corymbia</i> spp. and/or <i>Eucalyptus melanophloia</i> open forest to woodland on Cainozoic alluvial plains	11.3.19
Forb and/or grassland ± scattered <i>Atalaya hemiglauca</i> , <i>Flindersia maculosa</i> , <i>Acacia</i> spp. on alluvial plains	11.3.20
<i>Eucalyptus tereticornis</i> or <i>E. camaldulensis</i> woodland fringing drainage lines	11.3.25
<i>Eucalyptus moluccana</i> or <i>E. microcarpa</i> woodland to open forest on margins of alluvial plains	11.3.26
Freshwater wetlands	11.3.27
<i>Eucalyptus crebra</i> , <i>E. exserta</i> , <i>Melaleuca</i> spp. woodland on alluvial plains	11.3.29
<i>Eucalyptus crebra</i> , <i>Corymbia dallachiana</i> woodland on alluvial plains	11.3.30
<i>Ophiuros exaltatus</i> , <i>Dichanthium</i> spp. grassland on alluvial plains	11.3.31
<i>Allocasuarina luehmannii</i> open woodland on alluvial plains	11.3.32

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus platyphylla, Corymbia clarksoniana</i> woodland on alluvial plains	11.3.35
<i>Eucalyptus coolabah</i> fringing woodland on alluvial plains	11.3.37
<i>Eucalyptus melanophloia ± E. chloroclada</i> open woodland on undulating plains and valleys with sandy soils	11.3.39
<i>Dichanthium</i> spp., <i>Astrebla</i> spp. grassland on Cainozoic clay plains	11.4.4
<i>Eucalyptus orgadophila</i> open woodland on Cainozoic clay plains	11.4.13
<i>Eucalyptus crebra</i> and/or <i>E. populnea, Callitris glaucophylla, Angophora leiocarpa, Allocasuarina luehmannii</i> woodland on Cainozoic sand plains and/or remnant surfaces	11.5.1
<i>Eucalyptus crebra, Corymbia</i> spp., with <i>E. moluccana</i> on lower slopes of Cainozoic sand plains and/or remnant surfaces	11.5.2
<i>Eucalyptus populnea ± E. melanophloia ± Corymbia clarksoniana</i> on Cainozoic sand plains and/or remnant surfaces	11.5.3
<i>Eucalyptus chloroclada, Callitris glaucophylla, C. endlicheri, Angophora leiocarpa</i> woodland on Cainozoic sand plains and/or remnant surfaces	11.5.4
<i>Eucalyptus melanophloia</i> and/or <i>Callitris glaucophylla</i> woodland on Cainozoic sand plains and/or remnant surfaces, on deep red sands	11.5.5
<i>Eucalyptus acmenoides</i> and/or <i>Angophora leiocarpa</i> on Cainozoic sand plains and/or remnants	11.5.7

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca</i> spp., <i>Eucalyptus crebra</i> , <i>Corymbia intermedia</i> woodland on Cainozoic sand plains and/or remnant surfaces	11.5.8
<i>Eucalyptus crebra</i> and other <i>Eucalyptus</i> spp. and <i>Corymbia</i> spp. woodland on Cainozoic sand plains and/or remnant surfaces	11.5.9
<i>Corymbia clarksoniana</i> woodland and other <i>Corymbia</i> spp. and <i>Eucalyptus</i> spp. on Cainozoic sand plains and/or remnant surfaces	11.5.12
Semi-evergreen vine thicket on Cainozoic sand plains and/or remnant surfaces	11.5.15
<i>Eucalyptus moluccana</i> and/or <i>E. microcarpa</i> and/or <i>E. woollsiana</i> ± <i>E. crebra</i> woodland on Cainozoic sand plains	11.5.20
<i>Corymbia blossomei</i> ± <i>Callitris glaucocephala</i> ± <i>Eucalyptus crebra</i> ± <i>Angophora leiocarpa</i> woodland on Cainozoic sand plains and/or remnant surfaces	11.5.21
<i>Acacia harpophylla</i> and/or <i>Casuarina cristata</i> and <i>Eucalyptus thozetiana</i> or <i>E. microcarpa</i> woodland on lower scarp slopes on Cainozoic lateritic duricrust	11.7.1
<i>Acacia</i> spp. woodland on Cainozoic lateritic duricrust. Scarp retreat zone	11.7.2
<i>Eucalyptus persistens</i> , <i>Triodia mitchellii</i> open woodland on stripped margins of Cainozoic lateritic duricrust	11.7.3
<i>Eucalyptus decorticans</i> and/or <i>Eucalyptus</i> spp., <i>Corymbia</i> spp., <i>Acacia</i> spp., <i>Lysicarpus angustifolius</i> woodland on Cainozoic lateritic duricrust	11.7.4

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Shrubland on natural scalds on deeply weathered coarse-grained sedimentary rocks	11.7.5
<i>Corymbia citriodora</i> or <i>Eucalyptus crebra</i> woodland on Cainozoic lateritic duricrust	11.7.6
<i>Eucalyptus fibrosa</i> subsp. <i>nubila</i> ± <i>Corymbia</i> spp. ± <i>Eucalyptus</i> spp. woodland on Cainozoic lateritic duricrust	11.7.7
<i>Eucalyptus laevopinea</i> tall open forest on Cainozoic igneous rocks. Elevated plateaus	11.8.1
<i>Eucalyptus tereticornis</i> , <i>E. melliodora</i> woodland on Cainozoic igneous rocks	11.8.2
<i>Eucalyptus melanophloia</i> open woodland on Cainozoic igneous rocks	11.8.4
<i>Eucalyptus orgadophila</i> open woodland on Cainozoic igneous rocks	11.8.5
<i>Macropteranthes leichhardtii</i> thicket on Cainozoic igneous rocks	11.8.6
<i>Eucalyptus albens</i> , <i>E. crebra</i> woodland on Cainozoic igneous rocks	11.8.8
<i>Eucalyptus melanophloia</i> ± <i>E. orgadophila</i> woodland on fine-grained sedimentary rocks	11.9.2
<i>Dichanthium</i> spp., <i>Astrebla</i> spp. grassland on fine-grained sedimentary rocks	11.9.3
<i>Macropteranthes leichhardtii</i> thicket on fine-grained sedimentary rocks	11.9.8
<i>Eucalyptus crebra</i> woodland on fine-grained sedimentary rocks	11.9.9

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia citriodora</i> woodland on coarse-grained sedimentary rocks	11.10.1
<i>Acacia catenulata</i> or <i>A. shirleyi</i> open forest on coarse-grained sedimentary rocks. Crests and scarps	11.10.3
<i>Eucalyptus decorticans</i> , <i>Lysicarpus angustifolius</i> ± <i>Eucalyptus</i> spp., <i>Corymbia</i> spp., <i>Acacia</i> spp. woodland on coarse-grained sedimentary rocks	11.10.4
<i>Eucalyptus sphaerocarpa</i> ± <i>E. mensalis</i> , <i>E. saligna</i> tall open forest on coarse-grained sedimentary rocks. Tablelands	11.10.5
<i>Angophora leiocarpa</i> , <i>Callitris glaucophylla</i> open woodland on coarse-grained sedimentary rocks. Broad valleys	11.10.6
<i>Eucalyptus crebra</i> woodland on coarse-grained sedimentary rocks	11.10.7
<i>Callitris glaucophylla</i> woodland on coarse-grained sedimentary rocks	11.10.9
<i>Eucalyptus populnea</i> , <i>E. melanophloia</i> ± <i>Callitris glaucophylla</i> woodland on coarse-grained sedimentary rocks	11.10.11
<i>Eucalyptus populnea</i> woodland on medium to coarse-grained sedimentary rocks	11.10.12
<i>Eucalyptus</i> spp. and/or <i>Corymbia</i> spp. open forest on scarps and sandstone tablelands	11.10.13
<i>Eucalyptus crebra</i> ± <i>Acacia rhodoxylon</i> woodland on old sedimentary rocks with varying degrees of metamorphism and folding	11.11.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia shirleyi</i> or <i>A. catenulata</i> low open forest on old sedimentary rocks with varying degrees of metamorphism and folding	11.11.2
<i>Corymbia citriodora</i> , <i>Eucalyptus crebra</i> , <i>E. acmenoides</i> open forest on old sedimentary rocks with varying degrees of metamorphism and folding. Coastal ranges	11.11.3
<i>Eucalyptus crebra</i> woodland on old sedimentary rocks with varying degrees of metamorphism and folding. Coastal ranges	11.11.4
Microphyll vine forest ± <i>Araucaria cunninghamii</i> on old sedimentary rocks with varying degrees of metamorphism and folding	11.11.5
<i>Corymbia leichhardtii</i> , <i>C. clarksoniana</i> woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.6
<i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> , <i>Corymbia xanthope</i> woodland on serpentinite	11.11.7
<i>Eucalyptus shirleyi</i> woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.8
<i>Eucalyptus populnea</i> or <i>E. brownii</i> woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.9
<i>Eucalyptus persistens</i> low woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.12
<i>Eucalyptus crebra</i> woodland on deformed and metamorphosed sediments and interbedded volcanics	11.11.15

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus thozetiana</i> , <i>Acacia harpophylla</i> woodland on old sedimentary rocks with varying degrees of metamorphism and folding	11.11.19
<i>Eucalyptus platyphylla</i> woodland on old sedimentary rocks with varying degrees of metamorphism and folding. Lowlands	11.11.20
<i>Eucalyptus crebra</i> woodland on igneous rocks	11.12.1
<i>Eucalyptus melanophloia</i> woodland on igneous rocks	11.12.2
<i>Eucalyptus crebra</i> , <i>E. tereticornis</i> , <i>Angophora leiocarpa</i> woodland on igneous rocks especially granite	11.12.3
Semi-evergreen vine thicket and microphyll vine forest on igneous rocks	11.12.4
<i>Corymbia citriodora</i> open forest on igneous rocks (granite)	11.12.6
<i>Eucalyptus crebra</i> woodland with patches of semi-evergreen vine thicket on igneous rocks (boulder-strewn hillsides)	11.12.7
<i>Eucalyptus platyphylla</i> woodland on igneous rocks	11.12.9
<i>Eucalyptus crebra</i> , <i>Corymbia</i> spp., <i>E. acmenoides</i> woodland on igneous rocks. Coastal hills	11.12.13

Part 2

Cape York Peninsula Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Rhizophora stylosa</i> ± <i>Bruguiera gymnorhiza</i> closed forest, occurs as outer mangroves	3.1.1
<i>Avicennia marina</i> (grey mangrove) low open forest on landward side of tidal zone	3.1.2
<i>Ceriops tagal</i> ± <i>Avicennia marina</i> low closed forest on intertidal areas	3.1.3
<i>Sporobolus virginicus</i> closed-tussock grassland on coastal plains	3.1.5
Sparse hermland or bare saltpans on salt plains and saline flats	3.1.6
Semi-deciduous vine thicket to vine forest on beach dunes and ridges	3.2.2
<i>Acacia crassicarpa</i> on coastal dunes and beach ridges, woodland to open forest	3.2.5
<i>Corymbia intermedia</i> or <i>C. clarksoniana</i> woodland in wet coastal areas	3.2.7
<i>Eucalyptus tetrodonta</i> and <i>Corymbia clarksoniana</i> ± <i>E. brassiana</i> or <i>Erythrophleum chlorostachys</i> woodland on stabilised dunes	3.2.10
Low microphyll vine forest on coastal dunes and beach ridges	3.2.11
Araucarian microphyll vine forest on coastal dunefields and beach ridges	3.2.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca viridiflora</i> and <i>Neofabricia myrtifolia</i> low open woodland on beach ridges	3.2.15
<i>Asteromyrtus lysicephala</i> ± <i>Neofabricia myrtifolia</i> open heath on flat sand plains	3.2.18
<i>Melaleuca arcana</i> and <i>Thryptomene oligandra</i> open heath in swampy areas on sand plains	3.2.20
<i>Neofabricia myrtifolia</i> ± <i>Jacksonia thesioides</i> open to closed heath on dunefields	3.2.21
Sparse hermland and/or shrubland and bare sand areas predominantly on sand blows	3.2.26
Closed semi-deciduous mesophyll vine forest on loamy alluvia	3.3.1
Semi-deciduous mesophyll and/or notophyll vine forest on alluvia	3.3.2
Evergreen to semi-deciduous notophyll vine forest on alluvia on major watercourses	3.3.5
<i>Corymbia tessellaris</i> , <i>C. clarksoniana</i> open forest on coastal alluvial plains	3.3.8
<i>Lophostemon suaveolens</i> open forest on streamlines, swamps and alluvial terraces	3.3.9
<i>Melaleuca argentea</i> and/or <i>M. fluviatilis</i> ± <i>M. leucadendra</i> open forest fringing streams and creeks	3.3.10
<i>Melaleuca saligna</i> ± <i>M. viridiflora</i> , <i>Lophostemon suaveolens</i> woodland on drainage swamps	3.3.14
<i>Eucalyptus chlorophylla</i> ± <i>Corymbia clarksoniana</i> woodland on alluvial plains and colluvial fans	3.3.16

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia clarksoniana</i> and <i>Erythrophleum chlorostachys</i> woodland on alluvial plains	3.3.17
<i>Corymbia clarksoniana</i> ± <i>C. dallachiana</i> woodland on alluvial plains	3.3.18
<i>Corymbia clarksoniana</i> ± <i>C. dallachiana</i> woodland on floodplains	3.3.19
<i>Corymbia clarksoniana</i> ± <i>Erythrophleum chlorostachys</i> ± <i>Melaleuca viridiflora</i> woodland on alluvial plains	3.3.20
<i>Corymbia clarksoniana</i> ± <i>Syzygium eucalyptoides</i> woodland on lower slopes of sand ridges and in drainage depressions	3.3.21
<i>Corymbia clarksoniana</i> or <i>C. novoguineensis</i> woodland on alluvial plains	3.3.22
<i>Corymbia clarksoniana</i> or <i>C. polycarpa</i> woodland on stream levees	3.3.23
<i>Eucalyptus leptophleba</i> ± <i>Corymbia clarksoniana</i> woodland on alluvial deposits	3.3.24
<i>Eucalyptus leptophleba</i> ± <i>Corymbia tessellaris</i> ± <i>E. platyphylla</i> woodland on riverine levees and floodplains	3.3.25
<i>Corymbia nesophila</i> ± <i>Eucalyptus tetrodonta</i> woodland on sandstone footslopes and fans	3.3.26
<i>Corymbia nesophila</i> ± <i>Eucalyptus tetrodonta</i> ± <i>E. brassiana</i> woodland on alluvial sediments	3.3.27
<i>Eucalyptus platyphylla</i> ± <i>Corymbia clarksoniana</i> woodland on alluvial and colluvial plains	3.3.28

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia polycarpa</i> ± <i>C. curtipes</i> woodland on Mitchell River levees	3.3.29
<i>Corymbia tessellaris</i> ± <i>Eucalyptus acroleuca</i> woodland on levees	3.3.30
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia</i> spp. woodland on coastal plains	3.3.31
<i>Melaleuca viridiflora</i> ± <i>M. saligna</i> woodland in sinkholes and drainage depressions	3.3.32
<i>Thryptomene oligandra</i> and <i>Melaleuca viridiflora</i> woodland on sides of depressions	3.3.33
<i>Eucalyptus acroleuca</i> open woodland on floodplains in Lakefield National Park	3.3.35
<i>Eucalyptus chlorophylla</i> open woodland on alluvial plains in south of bioregion	3.3.36
<i>Eucalyptus microtheca</i> ± <i>Corymbia dallachiana</i> open woodland on floodplains	3.3.37
Deciduous notophyll and/or microphyll vine thicket ± <i>Lagerstroemia archeriana</i> on heavy clay alluvium	3.3.38
<i>Melaleuca clarksonii</i> low open forest in swamps	3.3.41
<i>Melaleuca viridiflora</i> low woodland in drainage areas	3.3.42
<i>Melaleuca viridiflora</i> ± <i>Xanthorrhoea johnsonii</i> low woodland on fans and alluvial plains	3.3.43
<i>Melaleuca citrolens</i> ± <i>M. foliolosa</i> low open woodland along drainage lines	3.3.47
<i>Melaleuca saligna</i> ± <i>M. viridiflora</i> low open woodland on drainage depressions and outwash plains	3.3.48

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca viridiflora</i> low open woodland on low plains	3.3.49
<i>Melaleuca viridiflora</i> ± <i>Petalostigma pubescens</i> ± <i>M. stenostachya</i> low open woodland on low plains	3.3.50
<i>Melaleuca citrolens</i> and/or <i>Excoecaria parvifolia</i> tall shrubland on eroding drainage areas	3.3.52
<i>Asteromyrtus lysicephala</i> ± <i>Baeckea frutescens</i> open heath on Jardine River sand plains	3.3.53
<i>Asteromyrtus lysicephala</i> and <i>Thryptomene oligandra</i> open heath on alluvial plains	3.3.55
<i>Eriachne</i> spp. ± <i>Aristida</i> spp. closed-tussock grassland on alluvial plains	3.3.56
<i>Oryza rufipogon</i> ± <i>Eleocharis</i> spp. closed-tussock grassland in seasonally inundated depressions	3.3.58
<i>Themeda arguens</i> , <i>Dichanthium sericeum</i> closed-tussock grassland on marine plains	3.3.60
<i>Panicum</i> spp. and <i>Fimbristylis</i> spp. tussock grassland on inland and coastal alluvial plains	3.3.61
<i>Eleocharis dulcis</i> dominated closed sedgeland on seasonally flooded marine plains	3.3.63
<i>Baloskion tetraphyllum</i> subsp. <i>meiostachyum</i> open sedgeland in drainage swamps in dune fields	3.3.64
Ephemeral lakes and lagoons on alluvial plains and depressions	3.3.65
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia stockeri</i> subsp. <i>peninsularis</i> tall woodland on deeply weathered plateaus	3.5.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tetrodonta</i> and <i>Corymbia nesophila</i> tall woodland on deeply weathered plateaus and remnants	3.5.2
Semi-deciduous notophyll vine forest in small patches on northern plateaus	3.5.4
<i>Eucalyptus phoenicea</i> ± <i>E. tetrodonta</i> woodland on sandy colluvia	3.5.6
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia clarksoniana</i> woodland on sand plains	3.5.7
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia stockeri</i> subsp. <i>peninsularis</i> ± <i>C. stockeri</i> woodland on erosional plains and sandstone plateaus	3.5.8
<i>Eucalyptus tetrodonta</i> and <i>Corymbia stockeri</i> subsp. <i>peninsularis</i> woodland on sand ridges	3.5.9
<i>Eucalyptus tetrodonta</i> and <i>Corymbia nesophila</i> woodland on sandy gently undulating rises and low hills	3.5.10
<i>Eucalyptus tetrodonta</i> and <i>Corymbia nesophila</i> woodland on lower slopes of plains and rises	3.5.11
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia nesophila</i> ± <i>C. clarksoniana</i> woodland on undulating rises	3.5.12
<i>Melaleuca viridiflora</i> ± <i>Acacia</i> spp. ± <i>Asteromyrtus symphyocarpa</i> low woodland on scattered coastal sand plains	3.5.14
<i>Melaleuca viridiflora</i> and <i>Asteromyrtus symphyocarpa</i> low woodland on colluvial plains	3.5.15
<i>Melaleuca viridiflora</i> ± <i>Neofabricia myrtifolia</i> low woodland on colluvial areas	3.5.16

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca viridiflora</i> and <i>M. stenostachya</i> low open woodland on flat plains	3.5.18
<i>Asteromyrtus lysicephala</i> and <i>Choriceras tricorne</i> open heath on sand sheets	3.5.19
<i>Corymbia clarksoniana</i> ± <i>Erythrophleum chlorostachys</i> ± <i>Corymbia</i> spp. woodland on plains	3.5.22
<i>Eucalyptus chlorophylla</i> ± <i>Corymbia clarksoniana</i> open woodland to woodland on undulating plains	3.5.24
<i>Eucalyptus leptophleba</i> ± <i>Corymbia tessellaris</i> , <i>E. platyphylla</i> or <i>C. clarksoniana</i> open woodland to woodland on undulating plains	3.5.25
<i>Eucalyptus platyphylla</i> ± <i>Corymbia clarksoniana</i> woodland to open forest on flat wet plains	3.5.26
<i>Melaleuca citrolens</i> ± <i>M. foliolosa</i> ± <i>M. viridiflora</i> low open woodland on plains	3.5.27
<i>Asteromyrtus lysicephala</i> ± <i>Thryptomene oligandra</i> ± <i>Neofabricia myrtifolia</i> open heath on pediment fans	3.5.28
<i>Sarga plumosum</i> ± <i>Themeda arguens</i> closed-tussock grassland on erosional plains	3.5.29
<i>Eucalyptus cullenii</i> ± <i>E. tetrodonta</i> woodland on erosional escarpments and plains	3.7.3
<i>Eucalyptus tetrodonta</i> and <i>Corymbia stockeri</i> subsp. <i>peninsularis</i> woodland on ironstone knolls and slopes	3.7.4
<i>Corymbia stockeri</i> , <i>Eucalyptus cullenii</i> woodland on ironstone knolls and erosional surfaces	3.7.5
<i>Melaleuca stenostachya</i> ± <i>Acacia leptostachya</i> woodland on lateritic erosional slopes	3.7.6

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia clarksoniana</i> ± <i>C. confertiflora</i> woodland on erosional plains	3.9.1
<i>Eucalyptus chlorophylla</i> woodland to open woodland on undulating clay plains	3.9.2
<i>Corymbia clarksoniana</i> woodland ± <i>Melaleuca viridiflora</i> on erosional plains	3.9.3
<i>Eucalyptus leptophleba</i> ± <i>Corymbia dallachiana</i> or <i>Eucalyptus platyphylla</i> open woodland on rolling plains	3.9.4
<i>Corymbia papuana</i> ± <i>Eucalyptus leptophleba</i> open woodland on rolling plains	3.9.5
<i>Piliostigma malabaricum</i> tall open shrubland on central clay plains	3.9.7
<i>Corymbia stockeri</i> subsp. <i>stockeri</i> ± <i>Eucalyptus tetrodonta</i> ± <i>E. cullenii</i> woodland on sandstone plateaus	3.10.6
<i>Eucalyptus phoenicea</i> ± <i>Corymbia nesophila</i> or <i>E. tetrodonta</i> woodland on wetter sandstone	3.10.7
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia stockeri</i> ± <i>C. clarksoniana</i> ± <i>C. tessellaris</i> woodland on sandstone plateaus	3.10.9
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia stockeri</i> sens. lat. woodland on sandstone plateaus and slopes	3.10.10
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia nesophila</i> woodland on undulating sandstone hills	3.10.11
<i>Asteromyrtus brassii</i> , <i>Neofabricia myrtifolia</i> and <i>Allocasuarina littoralis</i> low open forest on sandstone plains	3.10.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Neofabricia myrtifolia</i> and <i>Asteromyrtus brassii</i> low open forest on plains and low rises	3.10.13
<i>Eucalyptus chlorophylla</i> low open woodland on sandstone hillslopes	3.10.15
<i>Melaleuca stenostachya</i> ± <i>M. viridiflora</i> ± <i>M. citrolens</i> low open woodland on sandstone ranges	3.10.16
<i>Asteromyrtus lysicephala</i> ± <i>Jacksonia thesioides</i> open heath on undulating plains and slopes	3.10.18
<i>Asteromyrtus lysicephala</i> and <i>Neofabricia myrtifolia</i> dwarf open heath on sandstone plateaus and headlands	3.10.19
<i>Corymbia nesophila</i> ± <i>Eucalyptus crebra</i> or <i>E. tetrodonta</i> woodland to open forest on sandstone plateaus and slopes	3.10.21
Simple evergreen notophyll vine forest on exposed metamorphic and granitic slopes	3.11.3
<i>Eucalyptus cullenii</i> and <i>Corymbia clarksoniana</i> woodland on low metamorphic hills and rises	3.11.7
<i>Eucalyptus cullenii</i> ± <i>Corymbia clarksoniana</i> woodland on metamorphic ranges	3.11.8
<i>Eucalyptus cullenii</i> and <i>Corymbia stockeri</i> subsp. <i>peninsularis</i> woodland on metamorphic hills	3.11.9
<i>Corymbia stockeri</i> ± <i>Eucalyptus tetrodonta</i> or <i>E. crebra</i> woodland on metamorphic hills	3.11.10
<i>Corymbia stockeri</i> ± <i>Eucalyptus tetrodonta</i> woodland on hills and erosional surfaces	3.11.11
<i>Eucalyptus leptophleba</i> and <i>E. platyphylla</i> ± <i>Corymbia tessellaris</i> woodland on rolling metamorphic hills	3.11.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia nesophila</i> ± <i>E. brassiana</i> woodland on metamorphic hills and ranges	3.11.13
<i>Eucalyptus leptophleba</i> and <i>Corymbia dallachiana</i> ± <i>E. chlorophylla</i> or <i>Corymbia clarksoniana</i> open woodland on metamorphic hills	3.11.15
<i>Eucalyptus chlorophylla</i> ± <i>Melaleuca viridiflora</i> low open woodland to open woodland on metamorphic slopes	3.11.17
<i>Melaleuca stenostachya</i> ± <i>M. viridiflora</i> ± <i>M. citrolens</i> low open woodland on metamorphic footslopes	3.11.18
Notophyll vine forest on granitic slopes and plateaus	3.12.3
<i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> open forest on coastal ranges and lowlands	3.12.8
<i>Corymbia tessellaris</i> and <i>C. clarksoniana</i> open forest on coastal ranges	3.12.9
<i>Eucalyptus cullenii</i> ± <i>Corymbia clarksoniana</i> woodland on acid volcanic ranges	3.12.10
<i>Corymbia stockeri</i> subsp. <i>peninsularis</i> ± <i>Welchiodendron longivalve</i> woodland on ranges and hills	3.12.11
<i>Corymbia nesophila</i> ± <i>Eucalyptus crebra</i> woodland on wet coastal granitic hills	3.12.12
<i>Corymbia nesophila</i> and <i>C. stockeri</i> subsp. <i>peninsularis</i> woodland on acid volcanic hills	3.12.13
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia stockeri</i> subsp. <i>peninsularis</i> woodland on rises and ridges	3.12.14

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tetrodonta</i> ± <i>Corymbia nesophila</i> woodland on low hills on granites	3.12.15
<i>Melaleuca viridiflora</i> and <i>Asteromyrtus brassii</i> woodland on granitic hills	3.12.16
<i>Eucalyptus leptophleba</i> or <i>Corymbia clarksoniana</i> open woodland on igneous hills and ranges	3.12.17
<i>Eucalyptus leptophleba</i> and <i>Corymbia clarksoniana</i> woodland to open woodland on coastal hills	3.12.18
<i>Corymbia confertiflora</i> or <i>Eucalyptus tetrodonta</i> woodland on volcanic hills and ranges	3.12.19
Deciduous to semi-deciduous vine thicket to forest on granite slopes	3.12.21
<i>Melaleuca viridiflora</i> ± <i>Neofabricia myrtifolia</i> ± <i>Petalostigma pubescens</i> on low woodland on granitic ranges	3.12.26

Part 3 Central Queensland Coast Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Mangrove closed forest of marine clay plains and estuaries	8.1.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Samphire open forbland on saltpans and plains adjacent to mangroves	8.1.2
<i>Corymbia</i> spp. and/or <i>Eucalyptus</i> spp. and/or <i>Acacia</i> spp. and/or <i>Allocasuarina littoralis</i> low open forest on Pleistocene parabolic dunes	8.2.8
<i>Melaleuca leucadendra</i> and/or <i>M. fluviatilis</i> and/or <i>Casuarina cunninghamiana</i> ± <i>Syncarpia glomulifera</i> open forest on creek banks	8.3.3
Variable woodland to open forest, often including <i>Corymbia intermedia</i> , <i>Eucalyptus portuensis</i> , <i>C. trachyphloia</i> , <i>E. platyphylla</i> and <i>E. drepanophylla</i> on low hills on metamorphosed sediments	8.11.3
<i>Corymbia citriodora</i> or <i>Eucalyptus molluccana</i> woodland on metamorphosed sediments (subregion 4)	8.11.8
Evergreen notophyll feather palm vine forest of uplands and highlands on Mesozoic to Proterozoic igneous rocks	8.12.1
Evergreen notophyll to complex notophyll vine forest of uplands, highlands and foothills on Mesozoic to Proterozoic igneous rocks	8.12.2
Evergreen to semi-evergreen, notophyll to microphyll, vine forest to vine thicket of foothills and uplands on Mesozoic to Proterozoic igneous rocks	8.12.3
<i>Eucalyptus portuensis</i> and/or <i>Lophostemon confertus</i> and/or <i>E. exserta</i> and/or <i>Corymbia trachyphloia</i> and/or <i>E. fibrosa</i> open forest on Mesozoic to Proterozoic igneous rocks	8.12.5
<i>Eucalyptus drepanophylla</i> ± <i>E. platyphylla</i> ± <i>Corymbia clarksoniana</i> woodland on low to medium hills on Mesozoic to Proterozoic igneous rocks	8.12.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia citriodora</i> ± <i>Eucalyptus portuensis</i> ± <i>E. drepanophylla</i> (or <i>E. crebra</i>) open forest on hill slopes and undulating plateaus on Mesozoic to Proterozoic igneous rocks	8.12.7
<i>Eucalyptus tereticornis</i> ± <i>Corymbia intermedia</i> ± <i>Lophostemon suaveolens</i> woodland on undulating uplands on Mesozoic to Proterozoic igneous rocks	8.12.9
Semi-evergreen microphyll vine thicket ± <i>Araucaria cunninghamii</i> on islands and coastal headlands on Mesozoic to Proterozoic igneous rocks and Tertiary volcanics	8.12.11
<i>Eucalyptus tereticornis</i> and/or <i>Corymbia</i> spp. and/or <i>E. platyphylla</i> and/or <i>Lophostemon suaveolens</i> woodland to open forest on hill slopes on Mesozoic to Proterozoic igneous rocks	8.12.12
<i>Eucalyptus drepanophylla</i> and/or <i>E. crebra</i> and/or <i>E. exserta</i> and/or <i>Acacia spirorbis</i> and/or <i>Lophostemon confertus</i> low woodland on islands and headlands on Mesozoic to Proterozoic igneous rocks and Tertiary acid to intermediate volcanics	8.12.14
Semi-evergreen notophyll and/or microphyll to complex notophyll <i>Argyrodendron</i> spp. vine forest ± <i>Araucaria cunninghamii</i> of foothills and uplands on near-coastal ranges and islands on Mesozoic to Proterozoic igneous rocks	8.12.18
Semi-deciduous complex notophyll feather palm vine forest of sheltered gullies and slopes, of foothills and uplands on Mesozoic to Proterozoic igneous rocks	8.12.19

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus drepanophylla</i> and/or <i>E. platyphylla</i> ± <i>Corymbia</i> spp. ± <i>E. crebra</i> woodland on low gently undulating landscapes on Mesozoic to Proterozoic igneous rocks	8.12.20
<i>Eucalyptus drepanophylla</i> and/or <i>Corymbia clarksoniana</i> ± <i>C. erthrophloia</i> ± <i>E. platyphylla</i> ± <i>E. exserta</i> ± <i>C. trachyphloia</i> woodland on hills and ranges at low to moderate altitudes in drier areas	8.12.22
<i>Eucalyptus resinifera</i> and/or <i>E. portuensis</i> and/or <i>E. acmenoides</i> and/or <i>Allocasuarina</i> spp. closed forest on moist upper slopes of ranges on Mesozoic to Proterozoic igneous rocks	8.12.31
<i>Corymbia intermedia</i> ± <i>E. portuensis</i> ± <i>E. exserta</i> open forest to woodland with areas of <i>Allocasuarina</i> spp. ± <i>Banksia integrifolia</i> open forest on high ranges on Mesozoic to Proterozoic igneous rocks	8.12.32

Part 4

Channel Country Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> ± <i>Melaleuca</i> spp. woodland on levees and banks of major rivers	5.3.1
<i>Eucalyptus camaldulensis</i> ± <i>E. coolabah</i> open woodland on levees and banks of drainage lines	5.3.2

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> ± <i>Atalaya hemiglaucha</i> ± <i>Acacia cambagei</i> ± <i>Acacia georginae</i> ± <i>Acacia cyperophylla</i> woodland on drainage lines within ranges	5.3.4
<i>Eucalyptus coolabah</i> open woodland with <i>Muehlenbeckia florulenta</i> shrubland on braided channel systems	5.3.5
<i>Eucalyptus coolabah</i> open woodland on alluvial plains	5.3.6
<i>Eucalyptus coolabah</i> ± <i>Lysiphyllo gilvum</i> ± <i>Acacia cambagei</i> low open woodland on drainage lines	5.3.7
<i>Eucalyptus coolabah</i> low open woodland with <i>Muehlenbeckia florulenta</i> on braided drainage lines	5.3.8
<i>Acacia cambagei</i> ± <i>Eucalyptus coolabah</i> tall shrubland on braided channels	5.3.9
<i>Acacia cambagei</i> low open woodland ± <i>Senna artemisioides</i> subsp. <i>oligophylla</i> ± <i>Eremophila</i> spp. on alluvium	5.3.10
<i>Acacia georginae</i> tall shrubland with <i>Senna artemisioides</i> subsp. <i>oligophylla</i> ± <i>Eremophila freelingii</i> on alluvium	5.3.11
<i>Chenopodium auricomum</i> ± <i>Duma florulenta</i> open shrubland in swamps and some clay pans between dunes	5.3.12
<i>Muehlenbeckia florulenta</i> open shrubland on swamps	5.3.13
<i>Atriplex nummularia</i> open shrubland on clay pans between dunes	5.3.14
<i>Maireana</i> spp. open shrubland on clay pans between dunes and floodplains	5.3.15

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eragrostis australasica</i> open grassland on alluvial plains and clay pans between dunes	5.3.16
<i>Tecticornia</i> spp. open-succulent shrubland fringing playa lakes or clay pans	5.3.17
Short grasses ± forbs open hermland on braided channel systems	5.3.18
Seasonally variable sparse to open hermland on frequently flooded but not distinctly channelled areas on alluvial plains, common dominants include <i>Sporobolus mitchellii</i> and/or <i>Eragrostis setifolia</i> and/or a range of ephemeral herbs	5.3.19
<i>Eucalyptus coolabah</i> ± <i>Eucalyptus camaldulensis</i> open woodland fringing billabongs and permanent waterholes	5.3.20
Seasonally variable sparse to open hermland on infrequently flooded alluvia of major rivers, their distributaries and larger creeklines, <i>Atriplex</i> spp., <i>Sclerolaena</i> spp., <i>Astrebla</i> spp., <i>Asteraceae</i> spp. and/or short grasses	5.3.21
Sparse hermland on clay pans and lakes	5.3.22
<i>Acacia aneura</i> low woodland on Quaternary deposits	5.5.1
<i>Acacia aneura</i> ± <i>Acacia sibirica</i> ± <i>Eremophila latrobei</i> tall shrubland on Quaternary deposits	5.5.2
<i>Acacia aneura</i> , <i>Acacia sibirica</i> tall shrubland on Quaternary sand sheets	5.5.3
<i>Acacia sibirica</i> ± <i>Acacia aneura</i> ± <i>Eucalyptus</i> spp. open shrubland on Quaternary sediments	5.5.4

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia sibirica</i> ± <i>Eucalyptus</i> spp. open shrubland on crests and tops of sandstone ranges	5.5.5
<i>Archidendropsis basaltica</i> and/or <i>Acacia aneura</i> ± <i>Corymbia terminalis</i> low open woodland on sand plains	5.5.6
<i>Crotalaria eremaea</i> ± <i>Eragrostis eriopoda</i> open formland on isolated and/or deflated sand dunes on alluvium	5.6.1
<i>Acacia georginae</i> and/or <i>Acacia cambagei</i> , <i>Eremophila obovata</i> ± <i>Eremophila macdonnellii</i> tall shrubland on clay plains between sand dunes	5.6.2
<i>Atalaya hemiglaucha</i> ± <i>Acacia aneura</i> ± <i>Acacia</i> spp. ± <i>Corymbia terminalis</i> tall open shrubland on sand dunes	5.6.4
<i>Triodia basedowii</i> hummock grassland on sides of, or between, dunes	5.6.5
<i>Triodia basedowii</i> hummock grassland wooded with <i>Acacia</i> spp., <i>Senna</i> spp., <i>Grevillea</i> spp. ± <i>Eucalyptus</i> spp. on sand plains and dune fields	5.6.6
<i>Triodia basedowii</i> hummock grassland wooded with <i>Eucalyptus pachyphylla</i> on sand plains	5.6.7
<i>Zygochloa paradoxa</i> ± <i>Crotalaria eremaea</i> ± <i>Triodia basedowii</i> open grassland on sand dunes	5.6.8
<i>Acacia shirleyi</i> ± <i>Acacia catenulata</i> ± <i>Acacia aneura</i> ± <i>Acacia cyperophylla</i> tall shrubland on tops and scarps of residuals	5.7.1
<i>Acacia shirleyi</i> ± <i>Eucalyptus thozetiana</i> tall shrubland with <i>Triodia</i> spp. ± <i>Acacia aneura</i> ± <i>Acacia cyperophylla</i> on scarps of residuals	5.7.2

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus normantonensis</i> tall shrubland with <i>Triodia</i> spp. on slopes and plateau margins of residuals	5.7.3
<i>Eucalyptus thozetiana</i> tall shrubland with <i>Triodia</i> spp. ± <i>Eucalyptus normantonensis</i> on plateau margins and slopes of residuals	5.7.4
<i>Acacia sibirica</i> open shrubland with <i>Triodia</i> spp. ± <i>Acacia aneura</i> ± <i>Acacia shirleyi</i> open shrubland on crests and tops of ranges	5.7.5
<i>Acacia cambagei</i> tall shrubland with <i>Triodia</i> spp. ± <i>Senna</i> spp. on eroding pediments	5.7.6
<i>Acacia cambagei</i> tall shrubland with <i>Eragrostis xerophila</i> , <i>Sporobolus actinocladus</i> on sediments on undulating plains	5.7.7
<i>Aristida</i> spp., <i>Eriachne pulchella</i> open grassland wooded with <i>Eucalyptus</i> spp. ± <i>Acacia sibirica</i> on plains	5.7.9
<i>Aristida latifolia</i> and <i>Aristida contorta</i> sparse grassland wooded with <i>Acacia tetragonophylla</i> ± <i>Senna</i> spp. on weathered Cretaceous sediments	5.7.10
Fluctuating climax of <i>Atriplex</i> spp., <i>Sclerolaena</i> sp. ± short grasses, open herbland on mantled pediments with dense silcrete cover	5.7.11
<i>Acacia cyperophylla</i> ± <i>A. aneura</i> tall shrubland on scarpas and low hills	5.7.12
<i>Acacia cyperophylla</i> ± <i>A. cambagei</i> or <i>A. georginae</i> ± <i>Atalaya hemiglaucha</i> tall shrubland on drainage lines	5.7.13
<i>Acacia sibirica</i> , <i>Hakea eyreana</i> ± <i>Acacia aneura</i> ± <i>Eremophila freelingii</i> open shrubland on sandstones	5.7.14

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Senna</i> spp., <i>Eremophila</i> spp. ± <i>Acacia tetragonophylla</i> open shrubland on Tertiary limestone	5.9.1
<i>Senna artemisioides</i> subsp. <i>helmsii</i> ± <i>Senna artemisioides</i> subsp. <i>oligophylla</i> ± <i>Acacia georginae</i> ± <i>Acacia</i> spp. open shrubland on Cambrian limestone	5.9.2
<i>Astrebla</i> spp. hermland ± short grasses ± forbs on Cretaceous sediments	5.9.3
<i>Aristida contorta</i> ± short grasses ± forbs on Cretaceous sediments with dense gravel cover	5.9.4
<i>Atriplex</i> spp., <i>Sclerolaena</i> spp., <i>Salsola australis</i> open hermland on Cretaceous sediments	5.9.5

Part 5**Desert Uplands
Bioregion**

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia argyrodendron</i> low open woodland on alluvial plains (western)	10.3.1
<i>Acacia argyrodendron</i> with or without <i>Eucalyptus cambageana</i> open woodland on alluvial plains (eastern)	10.3.2
<i>Acacia harpophylla</i> and/or <i>Eucalyptus cambageana</i> low open woodland to open woodland on alluvial plains	10.3.3

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia cambagei</i> low open woodland to low woodland on alluvial plains	10.3.4
<i>Eucalyptus brownii</i> open woodland on alluvial plains	10.3.6
<i>Astrebla</i> spp., <i>Iseilema vaginiflorum</i> and/or <i>Dichanthium fecundum</i> or <i>Bothriochloa ewartiana</i> tussock grassland on alluvial plains	10.3.7
<i>Aristida latifolia</i> and <i>Brachyachne convergens</i> sparse-tussock grassland or <i>Sclerolaena</i> spp. dwarf open shrubland on alluvial plains	10.3.8
<i>Eucalyptus whitei</i> open woodland on sandy alluvial fans	10.3.9
<i>Corymbia dallachiana</i> and <i>C. terminalis</i> open woodland on old alluvial plains (western)	10.3.10
<i>Corymbia citriodora</i> or <i>C. leichhardtii</i> woodland to tall woodland on alluvium in valleys	10.3.11
<i>Corymbia dallachiana</i> and <i>C. plena</i> or <i>C. terminalis</i> open woodland on sandy alluvial terraces (eastern)	10.3.12
<i>Melaleuca fluviatilis</i> and/or <i>Eucalyptus camaldulensis</i> woodland along watercourses	10.3.13
<i>Eucalyptus camaldulensis</i> and/or <i>E. coolabah</i> open woodland along channels and on floodplains	10.3.14
Grasslands, sedgelands, ephemeral herblands and open woodland in depressions on sand plains	10.3.15
<i>Triodia longiceps</i> hummock grassland, ephemeral open herblands, and <i>Melaleuca bracteata</i> low woodland on alluvial plains	10.3.16

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Clay pans, <i>Fimbristylis</i> sp. (Lake Buchanan) open sedgeland and spare-tussock grasslands on shallow alluvial plains (Lake Buchanan)	10.3.22
<i>Halosarcia</i> spp. open-succulent shrubland, <i>Diplachne fusca</i> sparse-tussock grassland and bare clay pan on lake bed (Lake Galilee)	10.3.23
Ephemeral lake bed (Lake Buchanan)	10.3.24
<i>Eremophila mitchellii</i> tall open shrubland on alluvial plains	10.3.25
<i>Eucalyptus populnea</i> open woodland on alluvial plains	10.3.27
<i>Eucalyptus melanophloia</i> or <i>E. crebra</i> open woodland on sandy alluvial fans	10.3.28
<i>Acacia argyrodendron</i> open woodland on Cainozoic lake beds	10.4.1
<i>Acacia harpophylla</i> and/or <i>Eucalyptus cambageana</i> open woodland on Cainozoic lake beds	10.4.3
<i>Acacia cambagei</i> low woodland on Cainozoic lake beds	10.4.5
<i>Astrebla squarrosa</i> and <i>Iseilema vaginiflorum</i> ± <i>Dichanthium sericeum</i> and <i>Panicum laevinode</i> open-tussock grassland on Cainozoic lake beds	10.4.8
<i>Eucalyptus similis</i> and/or <i>Corymbia brachycarpa</i> and/or <i>Corymbia setosa</i> low open woodland on sand plains	10.5.1
<i>Corymbia plena</i> with or without <i>C. dallachiana</i> or <i>C. terminalis</i> woodland on sand plains	10.5.2

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> or <i>E. drepanophylla</i> open woodland on sand plains	10.5.4
<i>Eucalyptus melanophloia</i> open woodland on sand plains	10.5.5
<i>Grevillea striata</i> , <i>G. parallela</i> and <i>Acacia sericophylla</i> low open woodland or <i>Corymbia terminalis</i> open woodland on relict sand plains	10.5.7
<i>Corymbia setosa</i> with <i>Grevillea pteridifolia</i> and/or <i>Melaleuca nervosa</i> low open woodland on sand plains	10.5.8
<i>Corymbia leichhardtii</i> open woodland on sand plains	10.5.10
<i>Eucalyptus whitei</i> or <i>E. melanophloia</i> open woodland on red sand plateaus	10.5.11
<i>Eucalyptus populnea</i> open woodland on sand plains	10.5.12
<i>Eucalyptus whitei</i> open woodland or <i>Corymbia dallachiana</i> low open woodland or <i>Triodia pungens</i> open-hummock grassland on silcrete	10.7.1
<i>Eucalyptus persistens</i> or <i>Corymbia dallachiana</i> low open woodland or <i>Triodia pungens</i> hummock grassland on ferricrete above scarp	10.7.2
<i>Acacia shirleyi</i> woodland or <i>A. catenulata</i> low woodland at margins of plateaus	10.7.3
<i>Eucalyptus persistens</i> low open woodland on pediments below scarp	10.7.4
<i>Eucalyptus thozetiana</i> open woodland on scarp and on pediments below scarp	10.7.5
<i>Melaleuca</i> spp. and/or <i>Acacia leptostachya</i> shrubland on ferricrete (eastern)	10.7.7

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca</i> spp. and/or <i>Acacia</i> spp. open shrubland on ferricrete (western)	10.7.8
<i>Eucalyptus whitei</i> open woodland or <i>Corymbia setosa</i> low open woodland on ferricrete	10.7.10
<i>Eucalyptus melanophloia</i> low open woodland on ferricrete	10.7.11
<i>Eucalyptus drepanophylla</i> or <i>E. crebra</i> open woodland on ferricrete	10.7.12
Ephemeral sparse-tussock grassland ground below scarp	10.7.13
<i>Acacia argyrodendron</i> low open woodland or dwarf open shrubland of chenopods or scald on Cretaceous sediments	10.9.1
<i>Acacia cambagei</i> and/or <i>Eucalyptus thozetiana</i> low woodland to open woodland on calcareous sandstones	10.9.2
<i>Acacia harpophylla</i> and/or <i>Eucalyptus cambageana</i> open woodland to woodland on Mesozoic sediments	10.9.3
<i>Acacia cambagei</i> low woodland on Cretaceous sediments	10.9.6
<i>Acacia shirleyi</i> woodland or <i>A. catenulata</i> low open woodland on sandstone ranges	10.10.1
<i>Acacia burdekensis</i> or <i>A. julifera</i> low open woodland and bare rock platforms on sandstone ranges	10.10.2

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus exilipes</i> and/or <i>Corymbia leichhardtii</i> open woodland on sandstone ranges	10.10.4
<i>Corymbia trachyphloia</i> and/or <i>C. lamprophylla</i> or <i>Eucalyptus mediocris</i> open woodland on sandstone ranges	10.10.5

Part 6

Einasleigh Uplands Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> and/or <i>E. tereticornis</i> ± <i>Melaleuca</i> spp. ± <i>Casuarina cunninghamiana</i> fringing woodland on channels and levees	9.3.1
<i>Eucalyptus leptophleba</i> and/or <i>E. chlorophylla</i> ± <i>Corymbia dallachiana</i> woodland on river levees and terraces	9.3.2
<i>Corymbia</i> spp. and <i>Eucalyptus</i> spp. dominated mixed woodland on alluvial flats, levees and plains	9.3.3
<i>Eucalyptus brownii</i> ± <i>Eucalyptus</i> spp. ± <i>Corymbia</i> spp. open woodland on alluvial plains	9.3.5
<i>Eucalyptus platyphylla</i> ± <i>Eucalyptus</i> spp. ± <i>Corymbia</i> spp. woodland on alluvial plains	9.3.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Wetlands and seasonally inundated grasslands with a fringing open woodland of mixed <i>Eucalyptus</i> spp. on Tertiary surfaces	9.3.7
<i>Eucalyptus moluccana</i> woodland on alluvial deposits	9.3.8
<i>Melaleuca bracteata</i> low closed forest ± <i>Eucalyptus</i> spp. emergents or vine thicket species on swamps in basalt plains	9.3.10
Wetlands (sometimes ephemeral) with aquatic species and fringed with <i>Eucalyptus</i> spp. communities within basalt plains and flows	9.3.11
River beds and associated waterholes on major rivers and channels	9.3.12
<i>Melaleuca</i> spp., <i>Eucalyptus camaldulensis</i> and <i>Casuarina cunninghamiana</i> fringing open forest on streams and channels	9.3.13
<i>Melaleuca</i> spp. ± <i>Acacia</i> spp. ± <i>Syzygium</i> spp. ± <i>Leptospermum</i> spp. fringing woodland on channels and levees	9.3.14
<i>Eucalyptus tereticornis</i> ± <i>Casuarina cunninghamiana</i> ± <i>Melaleuca</i> spp. fringing woodland on channels and levees	9.3.15
<i>Eucalyptus tereticornis</i> ± <i>E. platyphylla</i> ± <i>Corymbia clarksoniana</i> woodland on alluvial flats, levees and plains	9.3.16
<i>Casuarina cunninghamiana</i> and/or <i>Eucalyptus camaldulensis</i> or <i>E. tereticornis</i> fringing open forest on channels and levees on basalt flows	9.3.17
<i>Eucalyptus coolabah</i> and/or <i>E. leptophleba</i> woodland on alluvial plains	9.3.19

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus microneura</i> ± <i>Corymbia</i> spp. ± <i>E. leptophleba</i> woodland on alluvial plains	9.3.20
<i>Eucalyptus crebra</i> or <i>E. cullenii</i> ± <i>Corymbia</i> spp. open woodland on alluvial levees and terraces	9.3.22
<i>Melaleuca viridiflora</i> and/or <i>M. citrolens</i> low woodland ± <i>Corymbia</i> spp. emergents on alluvial deposits	9.3.24
<i>Dichanthium</i> spp., and/or <i>Astrebla</i> spp. ± <i>Iseilema</i> spp. grassland on alluvial deposits derived from basalt soils	9.3.25
Mixed grassland to open grassland including <i>Eragrostis</i> sp., <i>Aristida</i> sp., <i>Enneapogon</i> sp., <i>Iseilema</i> sp., <i>Chloris</i> sp. or <i>Dichanthium</i> sp. on non-basalt derived alluvial deposits	9.3.26
<i>Iseilema</i> sp., <i>Dichanthium</i> sp. grassland ± <i>Eucalyptus</i> spp. or <i>Corymbia</i> spp. emergents on alluvials on basalt geologies	9.3.27
<i>Eucalyptus similis</i> open forest on red kandosols on Tertiary plateaus, mesas and tablelands	9.5.1
<i>Eucalyptus crebra</i> ± <i>E. drepanophylla</i> and <i>Corymbia clarksoniana</i> woodland on sand plains	9.5.3
<i>Eucalyptus melanophloia</i> ± <i>Corymbia dallachiana</i> woodland on sand plains	9.5.4
<i>Corymbia clarksoniana</i> , <i>Eucalyptus portuensis</i> , <i>E. crebra</i> and <i>C. citriodora</i> in mixed open forests on red kandosols on Tertiary surfaces	9.5.5
<i>Eucalyptus leptophleba</i> ± <i>Corymbia</i> spp. woodland on yellow kandosols on Tertiary remnant surfaces	9.5.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> and <i>Corymbia erythrophloia</i> ± <i>C. polycarpa</i> woodland on kandosols	9.5.7
<i>Eucalyptus cullenii</i> and/or <i>E. leptophleba</i> woodland on undulating plains on remnant Tertiary surfaces	9.5.8
<i>Corymbia clarksoniana</i> and/or <i>Eucalyptus leptophleba</i> and/or <i>E. platyphylla</i> woodland on plains	9.5.9
<i>Eucalyptus microneura</i> ± <i>Terminalia</i> spp. woodland on sand sheets	9.5.10
<i>Eucalyptus persistens</i> ± <i>E. crebra</i> woodland on flats on Tertiary remnant plains	9.5.11
<i>Eucalyptus chlorophylla</i> and/or <i>E. tardecidens</i> woodland on Tertiary plains	9.5.12
<i>Melaleuca citrolens</i> and/or <i>Macropteranthes montana</i> low woodland with <i>Eucalyptus</i> spp. emergents on Tertiary sand sheets	9.5.13
<i>Melaleuca monantha</i> ± <i>M. viridiflora</i> ± <i>Callitris intratropica</i> mixed low woodland on valley infill	9.5.15
<i>Eucalyptus tetrodonta</i> ± <i>Erythrophleum chlorostachys</i> woodland on Tertiary remnants	9.5.16
<i>Eucalyptus persistens</i> woodland on laterised and deeply weathered surfaces on undulating terrain	9.7.1
<i>Acacia shirleyi</i> low open forest to woodland on mesas and lateritised surfaces	9.7.2
<i>Eucalyptus crebra</i> or <i>E. portuensis</i> ± <i>Corymbia clarksoniana</i> woodland on lateritised surfaces and edges of Tertiary surfaces	9.7.3

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia setosa</i> and/or <i>C. peltata</i> low open woodland on lateritised and deeply weathered surfaces	9.7.5
<i>Eucalyptus crebra</i> ± <i>Corymbia dallachiana</i> ± <i>E. leptophleba</i> open woodland on plains and rocky rises of basalt geologies	9.8.1
<i>Eucalyptus leptophleba</i> ± <i>Corymbia clarksoniana</i> ± <i>C. erythrophloia</i> open woodland on basalt plains	9.8.2
<i>Eucalyptus crebra</i> ± <i>E. tereticornis</i> open woodland on basalt plains	9.8.4
<i>Astrebla</i> spp. ± <i>Iseilema vaginiflorum</i> grassland to <i>Corymbia terminalis</i> open woodland on basalt plains	9.8.5
<i>Acacia cambagei</i> low woodland on scree slopes and footslopes of basalt tablelands	9.8.6
Semi-evergreen vine thicket on cones, craters and rocky basalt flows with little soil development	9.8.7
<i>Eucalyptus orgadophila</i> ± <i>Corymbia</i> spp. open woodland on basalt plains and rocky basalt hills	9.8.9
<i>Eucalyptus microneura</i> ± <i>Corymbia</i> spp. ± <i>Terminalia</i> spp. woodland on basalt plains	9.8.11
<i>Iseilema</i> spp. and/or <i>Dichanthium</i> spp. tussock grassland on basalt plains	9.8.13
<i>Eucalyptus chartaboma</i> ± <i>E. tetrodonta</i> woodland on sandstone scarps and plateaus with shallow sandy soils	9.10.1
<i>Acacia shirleyi</i> ± mixed species woodland to open forest on sandstone	9.10.3
<i>Eucalyptus melanophloia</i> low woodland on skeletal soils on metamorphics hills	9.11.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> (or several other ironbark species) ± <i>Corymbia</i> spp. woodland on shallow texture contrast soils on low metamorphic hills and lowlands	9.11.2
<i>Eucalyptus cullenii</i> or <i>E. staigeriana</i> ± <i>Corymbia clarksoniana</i> woodland on skeletal soils on metamorphic hills	9.11.3
<i>Eucalyptus crebra</i> , <i>Corymbia clarksoniana</i> , <i>C. citriodora</i> ± <i>E. portuensis</i> open forest on shallow soils on metamorphic hills and ranges	9.11.4
<i>Eucalyptus persistens</i> ± <i>E. crebra</i> woodland on low metamorphic hills	9.11.5
<i>Eucalyptus platyphylla</i> and/or <i>E. cullenii</i> ± <i>Corymbia clarksoniana</i> woodland on texture contrast soils on metamorphic hills	9.11.7
Semi-evergreen vine thicket on limestone rock outcrops	9.11.8
<i>Eucalyptus cloeziana</i> , <i>Corymbia citriodora</i> , <i>E. portuensis</i> and <i>E. cullenii</i> mixed woodland on steep dissected hills on highly metalliferous metamorphic rocks	9.11.10
<i>Eucalyptus cullenii</i> , <i>Corymbia hylandii</i> and <i>E. tetrodonta</i> ± <i>Erythrophleum chlorostachys</i> woodland on metamorphic hills	9.11.12
<i>Eucalyptus cullenii</i> and <i>Corymbia hylandii</i> or <i>C. erythrophloia</i> open woodland on undulating plains and rises	9.11.13
<i>Eucalyptus crebra</i> and <i>Corymbia citriodora</i> ± <i>Corymbia</i> spp. woodland on metamorphic hills and mountains in far south-west of bioregion	9.11.14

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> and/or <i>E. cullenii</i> and/or <i>E. whitei</i> ± <i>Corymbia pocillum</i> or <i>C. erythrophloia</i> woodland on metamorphic hills	9.11.15
<i>Eucalyptus crebra</i> ± <i>Corymbia erythrophloia</i> or <i>C. pocillum</i> woodland on steep to rolling hills	9.11.16
<i>Corymbia peltata</i> ± <i>Eucalyptus crebra</i> ± <i>E. shirleyi</i> or <i>E. melanophloia</i> low open woodland on metamorphic hills and mountains	9.11.17
<i>Eucalyptus quadricostata</i> ± <i>Corymbia erythrophloia</i> ± <i>C. leichhardtii</i> open woodland on metamorphic hills and ranges	9.11.18
<i>Eucalyptus microneura</i> ± <i>Corymbia erythrophloia</i> or <i>C. pocillum</i> low open woodland on rolling metamorphic hills and rises	9.11.23
<i>Eucalyptus microneura</i> or <i>Melaleuca citrolens</i> or <i>E. whitei</i> low open woodland with <i>Triodia</i> spp. ground layer on metamorphic low gravelly hills and rises	9.11.24
<i>Eucalyptus tardecidens</i> or <i>E. chlorophylla</i> ± <i>Corymbia</i> spp. ± <i>E. cullenii</i> low woodland on steep to rolling metamorphic hills and rises	9.11.25
<i>Eucalyptus leptophleba</i> and/or <i>E. cullenii</i> and/or <i>Corymbia clarksoniana</i> ± <i>E. platyphylla</i> woodland on undulating terrain to rolling hills	9.11.26
<i>Acacia shirleyi</i> ± <i>Eucalyptus microneura</i> ± <i>E. crebra</i> woodland on metamorphic hills and outcrops	9.11.28
<i>Acacia leptostachya</i> low woodland with emergents on stony and rocky metamorphic hills	9.11.30

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia terminalis</i> and/or <i>Eucalyptus crebra</i> and/or <i>C. erythrophloia</i> woodland on aprons surrounding limestone outcrops	9.11.31
<i>Eucalyptus crebra</i> and/or <i>E. xanthoclada</i> and/or <i>E. drepanophylla</i> low open woodland on igneous rocks	9.12.1
<i>Eucalyptus portuensis</i> , <i>Corymbia citriodora</i> , <i>E. granitica</i> or <i>E. crebra</i> , <i>C. intermedia</i> or <i>C. clarksoniana</i> mixed woodland on steep hills and ranges on igneous hills close to Wet Tropics boundary	9.12.2
<i>Eucalyptus chartaboma</i> ± <i>Eucalyptus</i> spp. ± <i>Corymbia</i> spp. woodland on sandy soils on igneous rocks	9.12.3
<i>Eucalyptus shirleyi</i> and/or <i>E. melanophloia</i> and/or <i>Corymbia peltata</i> and/or <i>Callitris intratropica</i> low open woodland on igneous rocks	9.12.4
<i>Eucalyptus quadricostata</i> ± <i>C. peltata</i> open woodland on igneous hills and steep hills	9.12.5
<i>Eucalyptus microneura</i> ± <i>Terminalia</i> spp. ± <i>Corymbia</i> spp. low open woodland on igneous hills	9.12.6
<i>Eucalyptus cullenii</i> ± <i>Corymbia leichhardtii</i> ± <i>C. erythrophloia</i> woodland on igneous rocks	9.12.7
Semi-evergreen vine thicket on rocky outcrops and shallow soils of igneous rocks	9.12.8
<i>Eucalyptus crebra</i> and/or <i>E. whitei</i> ± <i>Corymbia erythrophloia</i> open woodland on steep to rolling hills on igneous rocks	9.12.11
<i>Eucalyptus crebra</i> and <i>Corymbia erythrophloia</i> ± <i>E. microneura</i> open woodland on igneous rocks	9.12.12

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus crebra</i> ± <i>Corymbia peltata</i> ± <i>C. pocillum</i> ± <i>Callitris intratropica</i> low woodland on hills and steep hills on igneous rocks	9.12.13
<i>Eucalyptus crebra</i> and <i>E. similis</i> ± <i>Callitris intratropica</i> low open woodland on hills on igneous rocks	9.12.14
<i>Eucalyptus staigeriana</i> ± <i>Erythrophleum chlorostachys</i> low open woodland on hills on igneous rocks	9.12.15
<i>Eucalyptus atrata</i> ± <i>Corymbia citriodora</i> ± <i>E. portuensis</i> woodland on mountains and hills on igneous rocks	9.12.17
<i>Eucalyptus crebra</i> or <i>E. exilipes</i> ± <i>Corymbia citriodora</i> ± <i>C. peltata</i> open woodland on granites with thin sand sheet	9.12.18
<i>Eucalyptus crebra</i> or <i>E. granitica</i> ± <i>Corymbia citriodora</i> ± <i>E. portuensis</i> mixed woodland on igneous hills	9.12.19
<i>Eucalyptus pachycalyx</i> ± <i>E. cloeziana</i> ± <i>Corymbia leichhardtii</i> woodland on steep igneous hills	9.12.20
<i>Eucalyptus drepanophylla</i> , <i>Corymbia clarksoniana</i> or <i>C. intermedia</i> and <i>C. dallachiana</i> woodland on steep rugged igneous ranges	9.12.22
<i>Eucalyptus drepanophylla</i> or <i>E. cebra</i> , <i>Corymbia leichhardtii</i> and <i>C. lampropyllya</i> low open woodland on igneous rocks	9.12.23
<i>Eucalyptus drepanophylla</i> or <i>E. crebra</i> and/or <i>E. xanthoclada</i> and <i>Corymbia peltata</i> woodland on igneous rocks	9.12.24

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus melanophloia</i> and/or <i>E. shirleyi</i> ± <i>Corymbia erythrophloia</i> low open woodland on igneous rocks	9.12.27
<i>Eucalyptus melanophloia</i> low open woodland, often with <i>E. crebra</i> , on low hills on igneous rocks	9.12.28
<i>Corymbia leichhardtii</i> and <i>Eucalyptus cloeziana</i> mixed woodland on igneous hills	9.12.30
<i>Eucalyptus persistens</i> woodland on rhyolites and granites	9.12.32
<i>Eucalyptus microneura</i> ± <i>Corymbia pocillum</i> low open woodland on igneous rocks	9.12.33
Semi-evergreen vine thicket with <i>Araucaria cunninghamii</i> on steep hills on igneous rocks	9.12.34
<i>Corymbia leichhardtii</i> , <i>C. lamprophylla</i> , <i>Pleiogynium timorense</i> ± <i>Araucaria cunninghamii</i> open woodland on igneous hills	9.12.35
<i>Cochlospermum gregorii</i> or <i>C. gillivraei</i> deciduous low woodland on rocky outcrops	9.12.36
<i>Acacia shirleyi</i> ± <i>Corymbia pocillum</i> ± <i>Eucalyptus microneura</i> woodland on igneous rocks	9.12.37
<i>Acacia shirleyi</i> ± <i>Eucalyptus shirleyi</i> woodland on igneous rocks	9.12.38
<i>Melaleuca citrolens</i> ± <i>Terminalia platyptera</i> ± <i>Corymbia dallachiana</i> low open woodland on rolling igneous hills	9.12.40
<i>Eucalyptus chlorophylla</i> low open woodland on undulating low granodiorite hills	9.12.44

[s 7]

Part 7 Gulf Plains Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Offshore tidal sands and mud flats, including sea grass beds	2.1.1
Tidal low coastal rises of shells, sand or mud, and associated gutters, usually with mangroves	2.1.2
Tidal channels and associated levees, usually with mangroves	2.1.3
Infrequently inundated clay plains and low samphire rises	2.1.4
Secondary dunes and swales	2.2.2
Grassland on low plains adjacent to estuarine zone	2.3.1
Freshwater and brackish wetlands in old river channels on low plains adjacent to estuarine zone	2.3.2
<i>Astrebla</i> spp. grassland on plains of cracking clays	2.3.3
<i>Dichanthium</i> spp. (blue grass) and <i>Eulalia aurea</i> (brown top) grassland on plains of cracking clays	2.3.4
<i>Lysiphyllo cunnighamii</i> woodland on plains of calcareous clays	2.3.5
Deciduous scrubs on plains of cracking clay	2.3.6
<i>Acacia cambagei</i> woodland on plains on clays	2.3.7
<i>Eucalyptus microtheca</i> , <i>Lysiphyllo cunnighamii</i> low open woodland and <i>Aristida</i> spp. on plains and low rises of texture contrast soils and earths	2.3.9

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus microtheca</i> and <i>Eucalyptus chlorophylla</i> low open woodland, and <i>Melaleuca viridiflora</i> woodlands and savannahs, on plains	2.3.10
<i>Eucalyptus microtheca</i> , <i>Excoecaria parvifolia</i> low open woodland and <i>Dichanthium</i> spp. on grey clay plains	2.3.11
<i>Eucalyptus microtheca</i> woodland to low open woodland with <i>Sorghum</i> spp. in seasonally flooded depressions on gleyed podsolics	2.3.15
Deepwater lagoons with water lilies and sedges	2.3.16
<i>Eucalyptus microtheca</i> woodland on channels in fine textured alluvial plains	2.3.17
<i>Atalaya hemiglaucha</i> and <i>Grevillea striata</i> low woodland on low rises and plains on red loamy soils	2.3.18
<i>Corymbia bella</i> , <i>Corymbia polycarpa</i> and <i>Eucalyptus pruinosa</i> woodland on low rises and plains on pale sandy soils	2.3.20
<i>Eucalyptus leptophleba</i> and <i>Corymbia</i> spp. woodland on low rises and plains on fine sands and red earths	2.3.21
<i>Corymbia polycarpa</i> and <i>Melaleuca</i> spp. woodland on sandy channels and levees	2.3.22
<i>Eucalyptus leptophleba</i> and <i>Corymbia confertiflora</i> woodland on sandy alluvial terraces and levees	2.3.23
<i>Melaleuca</i> spp. woodland to open forest on sands in channels and on levees	2.3.24
<i>Eucalyptus camaldulensis</i> woodland on levees and floodplains	2.3.25

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> and <i>Nauclea orientalis</i> open forest fringing major tributaries	2.3.26
<i>Eucalyptus leucophylla</i> and <i>Corymbia terminalis</i> woodland in depressions on podsolic soils	2.3.27
<i>Melaleuca</i> spp. woodland in depressions and shallow valleys on solodised soils and pale earths	2.3.28
<i>Melaleuca</i> spp. woodland fringing depressions and broad valleys on solodised soils	2.3.29
<i>Melaleuca</i> spp. woodland in seasonally flooded depressions on podsolic soils	2.3.30
<i>Melaleuca</i> spp. low woodland in depressions and valley bottoms on fine-textured yellow earths	2.3.31
<i>Aristida</i> spp. grassland in depressions and valley bottoms on fine-textured yellow earths	2.3.32
<i>Eucalyptus microtheca</i> open woodland and sedges in circular depressions in sand plains on cracking clays	2.3.33
<i>Eucalyptus camaldulensis</i> woodland and sedges in circular depressions on podsolic soils	2.3.34
<i>Melaleuca</i> spp. low woodland in bottoms of shallow valleys on solodised soils	2.3.36
<i>Lysiphylgium cunninghamii</i> , <i>Atalaya hemiglaucha</i> and <i>Grevillea striata</i> low woodland on plains on earths and sandy soils	2.5.1
<i>Atalaya hemiglaucha</i> and <i>Ventilago viminalis</i> low open woodland on plains on red and brown earths	2.5.2
Evergreen scrub on plains on mainly deep sandy soils	2.5.3

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tetrodonta</i> and <i>Corymbia polycarpa</i> open woodland on pale earths and sands on plains	2.5.5
<i>Eucalyptus tetrodonta</i> and <i>Corymbia</i> spp. woodland to open forest on plains on red and yellow earths	2.5.6
<i>Eucalyptus tetrodonta</i> open forest on plains on deep podsolic soils	2.5.8
<i>Eucalyptus microneura</i> woodland on plains and plateaus on earths, podsolics and skeletal soils	2.5.9
<i>Eucalyptus leucophylla</i> , <i>Corymbia terminalis</i> and <i>Eucalyptus tectifica</i> woodland on sand plains on podsolic soils	2.5.10
<i>Eucalyptus leucophloia</i> low open woodland on plains on lateritic podsolic soils	2.5.11
<i>Eucalyptus pruinosa</i> low woodland on plains and low rises on red and yellow earths	2.5.12
<i>Corymbia polycarpa</i> woodland on sand plains on lateritic podsolic soils	2.5.13
<i>Melaleuca</i> spp. low woodland on plains on earths and podsolics (south)	2.5.14
<i>Melaleuca</i> spp. woodland on plains on earths and podsolics (north)	2.5.15
<i>Melaleuca foliolosa</i> shrubland on dissected plains on alkaline earths and texture contrast soil	2.5.16
<i>Acacia shirleyi</i> low open forest or <i>Melaleuca tamariscina</i> shrubland on laterised mudstones on skeletal soils	2.7.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia shirleyi, Eucalyptus shirleyi, Corymbia setosa</i> subsp. <i>pedicellaris</i> or <i>Melaleuca acacioides</i> woodland on low scarps on skeletal soils	2.7.2
<i>Triodia</i> spp. grassland on plateaus on skeletal soils and shallow earths	2.7.3
<i>Eucalyptus leucophloia</i> low woodland on lateritic scarps on skeletal soils	2.7.4
<i>Terminalia canescens</i> and <i>Corymbia setosa</i> subsp. <i>pedicellaris</i> woodland on dissected plateau margins on skeletal soils	2.7.5
Eucalypt woodland on hills and lowlands on basalts	2.8.1
<i>Astrebla</i> spp. grassland downs on shales with cracking clay soils	2.9.1
<i>Dichanthium</i> spp., <i>Eulalia aurea</i> grassland on shales with cracking clay soils	2.9.2
Deciduous scrub and grasslands on deep cracking clays on mudstones	2.9.3
<i>Acacia cambagei</i> low woodland on shales with cracking clay soils	2.9.4
<i>Acacia cambagei</i> low woodland in depressions on sand plains	2.9.5
<i>Melaleuca</i> spp. and <i>Corymbia polycarpa</i> woodland on pale earths on mudstones	2.9.6
<i>Eucalyptus chlorophylla</i> woodland on lowlands on earths and clays	2.9.7
<i>Eucalyptus microneura</i> woodland on undulating plains on sandstones with sands and earths	2.10.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Mixed eucalypt woodland on plateaus, mesas and scarps on shallow soils	2.10.2
<i>Eucalyptus</i> spp., <i>Corymbia citriodora</i> and <i>Eucalyptus acmenoides</i> open forest on high plateaus on earths and sands	2.10.3
<i>Eucalyptus microneura</i> woodland and <i>Triodia pungens</i> hummock grassland on scarps and stony ledges	2.10.4
<i>Acacia shirleyi</i> woodland and <i>Triodia pungens</i> hummock grassland on scarps and stony ledges	2.10.5
<i>Melaleuca</i> spp. low open woodland on ledges on skeletal soils	2.10.6
Eucalypt woodland on Precambrian sandstones	2.10.7
Eucalypt woodland and deciduous woodland on stony hills on folded sediments	2.11.1
Eucalypt woodland and deciduous woodland on hills on granitic rocks	2.12.1

Part 8

Mitchell Grass Downs Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> ± <i>Melaleuca</i> spp. woodland on drainage lines	4.3.1
<i>Eucalyptus camaldulensis</i> ± <i>E. coolabah</i> woodland on drainage lines	4.3.2
<i>Eucalyptus coolabah</i> , <i>E. camaldulensis</i> ± <i>Lysiphylgium gilvum</i> open woodland on drainage lines	4.3.3
<i>Eucalyptus coolabah</i> open woodland on drainage lines and/or plains	4.3.4
<i>Eucalyptus coolabah</i> ± <i>E. camaldulensis</i> ± <i>Acacia georginae</i> open woodland on drainage lines and/or plains	4.3.5
<i>Atalaya hemiglaucha</i> ± <i>Acacia georginae</i> ± <i>A. cyperophylla</i> woodland on alluvium	4.3.6
<i>Acacia cambagei</i> low woodland on braided channels or alluvial plains	4.3.8
<i>Acacia georginae</i> and <i>Eragrostis setifolia</i> tall open shrubland on drainage lines and alluvial plains	4.3.9
<i>Corymbia terminalis</i> ± <i>Lysiphylgium gilvum</i> and <i>Acacia victoriae</i> low open woodland on alluvium	4.3.10
<i>Eucalyptus coolabah</i> ± <i>E. camaldulensis</i> open woodland on alluvium, billabongs and permanent waterholes	4.3.11
<i>Chenopodium auricomum</i> ± <i>Muehlenbeckia florulenta</i> open shrubland on swamps	4.3.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eragrostis setifolia</i> and <i>Marsilea drummondii</i> ± <i>Chenopodium auricomum</i> open grassland in drainage depressions	4.3.13
<i>Astrebla lappacea</i> , <i>Astrebla</i> spp. ± <i>Eulalia aurea</i> grassland on alluvium	4.3.14
<i>Astrebla squarrosa</i> ± <i>Dichanthium</i> spp. ± <i>Eulalia aurea</i> grassland on alluvium	4.3.15
<i>Astrebla elymoides</i> ± <i>A. squarrosa</i> ± <i>Aristida latifolia</i> grassland on alluvium	4.3.16
<i>Astrebla pectinata</i> ± <i>Astrebla</i> spp. ± <i>Aristida latifolia</i> grassland on alluvium	4.3.17
<i>Eulalia aurea</i> , <i>Astrebla squarrosa</i> ± <i>Astrebla</i> spp. grassland on alluvial plains	4.3.18
<i>Dichanthium</i> spp., <i>Eulalia aurea</i> , <i>Astrebla</i> spp. grassland on alluvium	4.3.19
<i>Atriplex</i> spp. and <i>Sclerolaena</i> spp. ± <i>Astrebla</i> spp. ± short grasses ± forbs open herbland on braided or flat alluvial plains	4.3.20
<i>Acacia tephrina</i> low woodland on alluvium	4.3.23
<i>Chenopodium auricomum</i> ± <i>Muehlenbeckia florulenta</i> open shrubland on depressions on floodplains	4.3.24
<i>Astrebla pectinata</i> ± <i>Aristida latifolia</i> ± <i>Eulalia aurea</i> grassland on Tertiary sediments overlying limestone	4.4.1
<i>Astrebla pectinata</i> and <i>Iseilema</i> spp. grassland	4.4.2
<i>Acacia aneura</i> ± <i>Atalaya hemiglaucha</i> ± <i>Grevillea striata</i> low woodland on sand plains	4.5.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia aneura</i> tall open shrubland on Quaternary sand sheets	4.5.2
<i>Acacia aneura</i> , <i>Triodia brizoides</i> or <i>Triodia molesta</i> tall open shrubland on Tertiary sand sheets	4.5.3
<i>Archidendropsis basaltica</i> and/or <i>Acacia aneura</i> ± <i>Corymbia terminalis</i> low open woodland on sand plains	4.5.4
<i>Corymbia terminalis</i> , <i>Triodia pungens</i> ± <i>Acacia</i> spp., <i>Senna</i> spp., <i>Eucalyptus</i> spp. low open woodland on sand plains	4.5.5
<i>Acacia cambagei</i> , <i>Senna</i> spp., <i>Sida platycalyx</i> tall open shrubland on Quaternary sand sheets	4.5.6
<i>Acacia georginae</i> (or <i>A. cambagei</i>), <i>Sida platycalyx</i> , <i>Sclerolaena cornishiana</i> tall open shrubland on Quaternary sand sheets	4.5.7
<i>Triodia pungens</i> hummock grassland wooded with <i>Acacia</i> spp. ± <i>Eucalyptus</i> spp. on Quaternary sand sheets	4.5.8
<i>Acacia cambagei</i> , <i>Archidendropsis basaltica</i> and mixed species open woodland on sand plains	4.5.9
<i>Acacia shirleyi</i> , <i>Triodia</i> spp. ± <i>Eucalyptus</i> spp. low woodland on scarp	4.7.1
<i>Eucalyptus normantonensis</i> tall open shrubland with <i>Triodia</i> spp. on plateau margins	4.7.2
<i>Acacia cambagei</i> tall open shrubland with <i>Triodia</i> spp. ± <i>Senna</i> spp. near eroding edges of Tertiary plateaus	4.7.4
<i>Acacia chisholmii</i> low shrubland	4.7.6

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus leucophylla</i> low open woodland ± <i>Corymbia terminalis</i> ± <i>Triodia</i> spp.	4.7.7
<i>Eucalyptus leucophloia</i> low open woodland	4.7.8
<i>Astrebla lappacea</i> ± <i>Aristida latifolia</i> ± <i>Panicum decompositum</i> grassland on Cretaceous sediments	4.9.1
<i>Astrebla lappacea</i> and <i>A. pectinata</i> ± <i>A. elymoides</i> grassland on Cretaceous sediments	4.9.2
<i>Astrebla pectinata</i> and herbs ± <i>Astrebla</i> spp. grassland on Cretaceous sediments	4.9.4
Seasonally variable mosaic of <i>Astrebla lappacea</i> and <i>Sclerolaena</i> spp. ± <i>Enneapogon</i> spp. open-tussock grassland and sparse to open forbland on Cretaceous sediments	4.9.5
<i>Astrebla</i> spp. grassland wooded with mixed tree species on Cretaceous sediments	4.9.6
<i>Astrebla</i> spp. grassland wooded with <i>Acacia tephrina</i> ± <i>A. cambagei</i> and <i>Atalaya hemiglaucha</i> on Cretaceous sediments	4.9.7
<i>Astrebla</i> spp. grassland wooded with <i>Atalaya hemiglaucha</i> ± <i>Alectryon oleifolius</i> ± <i>Flindersia maculosa</i> on Cretaceous sediments	4.9.8
<i>Astrebla</i> spp. grassland wooded with <i>Acacia sutherlandii</i> or <i>A. victoriae</i> on Cretaceous sediments	4.9.9
<i>Acacia georginae</i> tall open shrubland on Cambrian limestone	4.9.10
<i>Corymbia terminalis</i> low open woodland with <i>Astrebla pectinata</i> ± <i>Eulalia aurea</i> on plains and low lying areas	4.9.12

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Senna artemisioides</i> subsp. <i>helmsii</i> ± <i>S. artemisioides</i> subsp. <i>oligophylla</i> ± <i>Acacia georginae</i> ± <i>Acacia</i> spp. open shrubland on tops and footslopes of Cambrian limestone residuals	4.9.13
<i>Acacia georginae</i> or <i>A. cambagei</i> low open woodland with <i>Astrebla</i> spp. on limestone	4.9.14
<i>Acacia cambagei</i> ± scattered shrub species including <i>Santalum lanceolatum</i> and <i>Eremophila mitchellii</i> tall open shrubland	4.9.16
<i>Archidendropsis basaltica</i> and mixed species including <i>Ventilago viminalis</i> and <i>Lysiphyllo caronii</i> on Cretaceous sediments	4.9.18
Clumps of <i>Acacia harpophylla</i> low woodland to tall shrubland with <i>Astrebla</i> spp. grassland on Cretaceous sediments sometimes with a covering of Tertiary deposits	4.9.19
<i>Astrebla lappacea</i> ± <i>Aristida latifolia</i> ± <i>Panicum decompositum</i> grassland on Cretaceous sediments with ashy soils	4.9.20

Part 9 **Mulga Lands Bioregion**

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus camaldulensis</i> woodland on alluvium within <i>Acacia aneura</i> associations	6.3.1
<i>Eucalyptus camaldulensis</i> ± <i>E. coolabah</i> ± <i>Acacia cambagei</i> woodland on major drainage lines or rivers	6.3.2
<i>Eucalyptus camaldulensis</i> ± <i>E. coolabah</i> ± <i>E. populnea</i> , <i>Acacia stenophylla</i> woodland on alluvium	6.3.3
<i>Acacia cambagei</i> ± <i>Eucalyptus ochrophloia</i> woodland on alluvium	6.3.4
<i>Eucalyptus ochrophloia</i> ± <i>Acacia cambagei</i> ± <i>E. coolabah</i> woodland on alluvium	6.3.5
<i>Acacia cambagei</i> low woodland on braided channels or alluvial plains	6.3.6
<i>Eucalyptus coolabah</i> , <i>Acacia stenophylla</i> low open woodland on alluvium	6.3.7
<i>Eucalyptus largiflorens</i> ± <i>Acacia cambagei</i> woodland on alluvium	6.3.8
<i>Eucalyptus coolabah</i> , <i>E. populnea</i> open woodland on alluvium	6.3.9
<i>Tecticornia</i> spp. open-succulent shrubland on alluvium	6.3.10
<i>Eleocharis pallens</i> ± short grasses ± <i>Eragrostis australasica</i> open herbland on clays, associated with ephemeral lakes, billabongs and permanent waterholes	6.3.11
<i>Acacia omalophylla</i> ± <i>A. microsperma</i> ± <i>Eucalyptus coolabah</i> tall open shrubland on alluvium	6.3.12

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Atriplex</i> spp., <i>Sclerolaena</i> spp., species of Asteraceae and/or short grasses open hermland on alluvial plains	6.3.13
<i>Astrebla</i> spp., <i>Dichanthium</i> spp. open grassland on alluvium	6.3.14
<i>Astrebla lappacea</i> , <i>A. pectinata</i> ± <i>A. elymoides</i> grassland on alluvium	6.3.15
<i>Callitris glauophylla</i> , <i>Acacia excelsa</i> , <i>Geijera parviflora</i> ± <i>Acacia aneura</i> woodland on alluvial dunes	6.3.16
<i>Callitris glauophylla</i> , <i>Corymbia tessellaris</i> , <i>Acacia excelsa</i> ± <i>C. clarksoniana</i> open woodland on old alluvial dunes and sand plains	6.3.17
<i>Eucalyptus populnea</i> ± <i>Eremophila mitchellii</i> ± <i>Acacia aneura</i> ± <i>E. melanophloia</i> woodland on flat alluvial plains	6.3.18
<i>Acacia aneura</i> , <i>A. excelsa</i> and/or <i>Geijera parviflora</i> low woodland on low alluvial sand dunes	6.3.21
<i>Acacia victoriae</i> ± <i>Eucalyptus</i> spp. tall open shrubland on old levees	6.3.22
<i>Eucalyptus coolabah</i> or <i>E. populnea</i> woodland on alluvial plains	6.3.24
<i>Acacia harpophylla</i> and/or <i>A. cambagei</i> low woodland to woodland on alluvial plains	6.3.25
<i>Acacia harpophylla</i> and/or <i>A. cambagei</i> low woodland on Quaternary deposits overlying older sediments	6.4.4
<i>Acacia aneura</i> , <i>Eucalyptus populnea</i> , <i>E. melanophloia</i> open forest on undulating lowlands	6.5.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus populnea</i> , <i>Acacia aneura</i> and/or <i>E. melanophloia</i> woodland on Quaternary sediments	6.5.2
<i>Acacia aneura</i> , <i>Eucalyptus populnea</i> low woodland on run-on plains	6.5.6
<i>Acacia aneura</i> , <i>Eucalyptus populnea</i> ± <i>E. intertexta</i> low woodland on run-on areas	6.5.7
<i>Acacia aneura</i> , <i>Eucalyptus populnea</i> ± <i>Eremophila gilesii</i> low woodland	6.5.8
<i>Acacia aneura</i> , <i>Eucalyptus populnea</i> ± <i>E. melanophloia</i> shrubby low woodland on Quaternary sediments	6.5.9
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> ± <i>Grevillea striata</i> , <i>A. excelsa</i> , <i>Hakea ivoryi</i> low woodland on sand plains	6.5.10
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> low woodland on sand plains	6.5.11
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> ± <i>E. melanophloia</i> ± <i>Brachychiton populneus</i> low woodland on sand plains	6.5.13
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> ± <i>Eremophila gilesii</i> tall open shrubland on Quaternary sediments	6.5.14
<i>Acacia aneura</i> , <i>Eucalyptus populnea</i> ± <i>Eremophila sturtii</i> tall open shrubland on sand plains	6.5.15
<i>Acacia aneura</i> groved with <i>Corymbia terminalis</i> or <i>C. blakei</i> tall open shrubland on Quaternary sediments	6.5.16
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> ± <i>E. melanophloia</i> ± <i>Eremophila mitchellii</i> low open woodland on plains	6.5.18

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Callitris glauophylla</i> ± <i>Angophora melanoxylon</i> ± <i>Eucalyptus melanophloia</i> ± <i>E. chloroclada</i> open woodland on Cainozoic sediments derived from old alluvial levees and dunes	6.5.19
<i>Atalaya hemiglaucha</i> ± <i>Acacia aneura</i> ± <i>Acacia</i> spp. ± <i>Corymbia terminalis</i> tall open shrubland on low dunes over alluvium	6.6.1
<i>Triodia mitchellii</i> ± <i>T. marginata</i> hummock grassland wooded with <i>Eucalyptus melanophloia</i> ± <i>Eucalyptus</i> spp. and <i>Acacia</i> spp. on low dunes	6.6.2
<i>Acacia catenulata</i> ± <i>A. shirleyi</i> ± <i>Eucalyptus</i> spp. open scrub on crests and slopes	6.7.1
<i>Acacia microsperma</i> open forest on upper and footslopes	6.7.2
<i>Eucalyptus thozetiana</i> or <i>E. cambageana</i> , <i>Acacia harpophylla</i> woodland on scarp	6.7.5
<i>Eucalyptus thozetiana</i> ± <i>Acacia aneura</i> open woodland on scarp and slopes	6.7.6
<i>Acacia catenulata</i> ± <i>Eucalyptus thozetiana</i> and/or <i>A. ensifolia</i> low open woodland with <i>Triodia</i> spp. and/or <i>A. petraea</i> ± <i>A. aneura</i> on scarp and plateaus	6.7.7
<i>Acacia aneura</i> ± <i>A. clivicola</i> ± <i>Eremophila latrobei</i> tall open shrubland on residuals	6.7.9
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> ± <i>Corymbia terminalis</i> tall shrubland on residuals	6.7.10
<i>Acacia aneura</i> ± <i>Eucalyptus cambageana</i> ± <i>Eucalyptus thozetiana</i> ± <i>Eremophila latrobei</i> tall shrubland on residuals	6.7.11

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia aneura</i> ± <i>Eucalyptus populnea</i> ± <i>E. melanophloia</i> ± <i>Eremophila gilesii</i> tall shrubland on residuals	6.7.12
<i>Acacia catenulata</i> ± <i>A. petraea</i> tall shrubland on scarp and tops of ranges	6.7.13
<i>Acacia clivicola</i> ± <i>Eucalyptus</i> spp. open shrubland on crests and tops of residuals	6.7.14
<i>Acacia brachystachya</i> , <i>A. aneura</i> open shrubland on the lower slopes of residuals	6.7.15
<i>Acacia clivicola</i> , <i>Eucalyptus exserta</i> open shrubland on colluvials associated with residuals	6.7.16
<i>Eriachne mucronata</i> open grassland wooded with <i>Acacia aneura</i> and/or <i>Corymbia terminalis</i> on plains or flat tops of residuals	6.7.17
<i>Acacia tephrina</i> ± <i>A. cambagei</i> low open woodland on undulating plains over Cretaceous sediments	6.9.2
<i>Acacia harpophylla</i> woodland with emergent <i>Eucalyptus cambageana</i> with stony soils derived from Cretaceous sediments	6.9.3
<i>Acacia cambagei</i> , <i>Senna</i> spp., <i>Sida platycalyx</i> tall open shrubland on undulating mantled pediments and scarp retreat zones	6.9.4

[s 7]

Part 10 New England Tableland Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus youmanii, E. dealbata, E. caleyi, Callitris endlicheri</i> woodland on metamorphics	13.11.1
<i>Eucalyptus melanophloia</i> woodland on metamorphics	13.11.4
<i>Corymbia citriodora</i> subsp. <i>variegata</i> open forest on metamorphics	13.11.6
<i>Eucalyptus campanulata</i> open forest on igneous rocks	13.12.1
<i>Eucalyptus andrewsii, E. youmanii</i> woodland on igneous rocks	13.12.2
<i>Eucalyptus youmanii</i> on igneous rocks	13.12.5

Part 11 Northwest Highlands Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Astrebla</i> spp. grassland on alluvial plains	1.3.1
<i>Eucalyptus microtheca</i> low open woodland to woodland on alluvial floodplains and channels	1.3.2

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Acacia cambagei</i> low open woodland to woodland on earths in valleys	1.3.4
Mixed eucalypt open woodland on sandy alluvial terraces	1.3.5
<i>Corymbia aparrerinja, Corymbia terminalis</i> open woodland on sandy terraces	1.3.6
<i>Eucalyptus camaldulensis</i> woodland on channels and levees (south)	1.3.7
<i>Eucalyptus camaldulensis</i> woodland on channels and levees (north)	1.3.8
<i>Eucalyptus leucophloia</i> low open woodland on red earths on plateaus	1.5.3
<i>Eucalyptus leucophylla</i> low open woodland on red earths in valleys	1.5.4
<i>Eucalyptus pruinosa</i> low open woodland on red earth plains	1.5.5
<i>Atalaya hemiglaucha, Ventilago viminalis, Grevillea striata</i> low open woodland on red earth plains	1.5.6
<i>Corymbia terminalis</i> and/or <i>Acacia aneura</i> low open woodland on sandy red earth plains	1.5.7
<i>Acacia cambagei</i> and <i>Atalaya hemiglaucha</i> low open woodland on red earth plains	1.5.8
<i>Ventilago viminalis</i> low open woodland on loams on sand sheet margins	1.5.9
<i>Eucalyptus leucophloia</i> low open woodland on skeletal soils on lateritic scarp and plateaus	1.7.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus pruinosa</i> low open woodland on calcareous red/brown earths	1.7.2
<i>Astrebla</i> spp. (Mitchell grass) grassland on shallow clays on limestones	1.9.1
Mixed shrubby woodland on rocky limestone hills	1.9.4
<i>Eucalyptus leucophylla</i> low open woodland to woodland on low hills on limestones and calcareous shales	1.9.5
<i>Eucalyptus leucophloia</i> and <i>Corymbia terminalis</i> low open woodland on limestone hills	1.9.6
<i>Eucalyptus pruinosa</i> low open woodland on shale hills	1.9.7
<i>Corymbia capricornia</i> low open woodland on sandstone plateaus	1.10.1
<i>Eucalyptus miniata</i> woodland on sandstone plateaus	1.10.2
<i>Corymbia aspera</i> low open woodland on rocky soils	1.10.3
<i>Eucalyptus leucophloia</i> and/or <i>Acacia</i> spp. low open woodland on stony sandstone plateaus	1.10.4
<i>Acacia shirleyi</i> open forest on skeletal soils and earths on sandstone plateaus	1.10.5
<i>Eucalyptus leucophloia</i> and <i>Corymbia grandifolia</i> low open woodland on stony low hills and colluvium	1.10.7
<i>Eucalyptus pruinosa</i> open woodland on slopes adjoining sandstone plateaus	1.10.8
Mixed shrubby woodland on folded limestones	1.11.1
<i>Eucalyptus leucophloia</i> low open woodland	1.11.2

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Corymbia terminalis</i> low open woodland on basic metamorphics	1.11.3
<i>Eucalyptus pruinosa</i> low open woodland on shallow soils in valleys below folded sediments	1.11.4
<i>Eucalyptus leucophloia</i> low open woodland on granites	1.12.1
<i>Eucalyptus melanophloia</i> (silver-leaved ironbark) low open woodland on low hills and torfields on biotite granites	1.12.2

Part 12 Southeast Queensland Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Saltpan vegetation including grassland, hermland and sedgeland on marine clay plains	12.1.2
Mangrove shrubland to low closed forest on marine clay plains and estuaries	12.1.3
<i>Corymbia intermedia</i> ± <i>Lophostemon confertus</i> ± <i>Banksia</i> spp. ± <i>Callitris columellaris</i> open forest on beach ridges usually in southern half of bioregion	12.2.5
<i>Eucalyptus racemosa</i> open forest on dunes and sand plains, usually deeply leached soils	12.2.6

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Melaleuca quinquenervia</i> or, rarely, <i>M. dealbata</i> open forest on sand plains	12.2.7
<i>Eucalyptus pilularis</i> open forest on parabolic high dunes	12.2.8
<i>Banksia aemula</i> low open woodland on dunes and sand plains, usually deeply leached soils	12.2.9
Mallee <i>Eucalyptus planchoniana</i> ± <i>Corymbia gummiifera</i> , <i>E. racemosa</i> subsp. <i>racemosa</i> , <i>Banksia aemula</i> woodland on dunes and sand plains, especially southern sand mass islands, usually deeply leached soils	12.2.10
<i>Corymbia tessellaris</i> ± <i>Eucalyptus tereticornis</i> , <i>C. intermedia</i> and <i>Livistona decora</i> woodland on beach ridges in northern half of bioregion	12.2.11
Closed heath on seasonally waterlogged sand plains	12.2.12
Foredune complex	12.2.14
<i>Gahnia sieberiana</i> , <i>Empodisma minus</i> , <i>Gleichenia</i> spp. closed sedgeland in coastal swamps	12.2.15
<i>Melaleuca quinquenervia</i> open forest on coastal alluvium	12.3.5
<i>Melaleuca quinquenervia</i> ± <i>Eucalyptus tereticornis</i> , <i>Lophostemon suaveolens</i> open forest on coastal alluvial plains	12.3.6
<i>Eucalyptus tereticornis</i> , <i>Casuarina cunninghamiana</i> subsp. <i>cunninghamiana</i> ± <i>Melaleuca</i> spp. fringing woodland	12.3.7
<i>Eucalyptus latisinensis</i> or <i>E. exserta</i> , <i>Melaleuca viridiflora</i> var. <i>viridiflora</i> woodland on alluvial plains	12.3.12

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Closed heathland on seasonally waterlogged alluvial plains usually near coast	12.3.13
Open forest complex with <i>Corymbia citriodora</i> subsp. <i>variegata</i> on subcoastal remnant Tertiary surfaces, usually deep red soils	12.5.1
<i>Eucalyptus latisinensis</i> ± <i>Corymbia intermedia</i> , <i>C. trachyphloia</i> subsp. <i>trachyphloia</i> , <i>Angophora leiocarpa</i> , <i>Eucalyptus exserta</i> woodland on complex of remnant Tertiary surfaces and Cainozoic and Mesozoic sediments	12.5.4
<i>Corymbia citriodora</i> subsp. <i>variegata</i> ± <i>Eucalyptus portuensis</i> or <i>E. acmenoides</i> , <i>E. fibrosa</i> subsp. <i>fibrosa</i> open forest on remnant Tertiary surfaces, usually deep red soils	12.5.7
<i>Eucalyptus latisinensis</i> and/or <i>Banksia aemula</i> low open woodland on complex of remnant Tertiary surface and Tertiary sedimentary rocks	12.5.10
<i>Eucalyptus campanulata</i> tall open forest on Cainozoic igneous rocks	12.8.1
Complex notophyll vine forest on Cainozoic igneous rocks at altitude of less than 600m	12.8.3
Complex notophyll vine forest with <i>Araucaria</i> spp. on Cainozoic igneous rocks	12.8.4
Complex notophyll vine forest on Cainozoic igneous rocks, usually at altitude of more than 600m	12.8.5
<i>Lophostemon confertus</i> open forest on Cainozoic igneous rocks	12.8.9

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus eugenoides</i> , <i>E. biturbinata</i> , <i>E. melliodora</i> ± <i>E. tereticornis</i> , <i>Corymbia intermedia</i> woodland on Cainozoic igneous rocks	12.8.14
<i>Eucalyptus melanophloia</i> ± <i>E. crebra</i> , <i>E. tereticornis</i> , <i>Corymbia tessellaris</i> woodland on Cainozoic igneous rocks	12.8.17
<i>Corymbia citriodora</i> subsp. <i>variegata</i> ± <i>Eucalyptus crebra</i> open forest on sedimentary rocks	12.9–10.2
<i>Eucalyptus racemosa</i> subsp. <i>racemosa</i> woodland on sedimentary rocks	12.9–10.4
Woodland complex often with <i>Corymbia trachyphloia</i> subsp. <i>Trachyphloia</i> , <i>C. citriodora</i> subsp. <i>variegata</i> , <i>Eucalyptus crebra</i> , <i>E. fibrosa</i> subsp. <i>fibrosa</i> on quartzose sandstone	12.9–10.5
<i>Eucalyptus pilularis</i> tall open forest on sedimentary rocks	12.9–10.14
<i>Eucalyptus acmenoides</i> , <i>E. major</i> , <i>E. siderophloia</i> ± <i>Corymbia citriodora</i> subsp. <i>variegata</i> woodland on sedimentary rocks	12.9–10.17
<i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> woodland on sedimentary rocks	12.9–10.19
<i>Eucalyptus acmenoides</i> or <i>E. portuensis</i> woodland, usually with <i>Corymbia trachyphloia</i> subsp. <i>trachyphloia</i> , on Cainozoic to Proterozoic sediments	12.9–10.21
Simple notophyll vine forest often with abundant <i>Archontophoenix cunninghamiana</i> (gully vine forest) on metamorphics ± interbedded volcanics	12.11.1

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus saligna</i> or <i>E. grandis</i> , <i>E. microcorys</i> , <i>Lophostemon confertus</i> tall open forest on metamorphics ± interbedded volcanics	12.11.2
<i>Eucalyptus siderophloia</i> , <i>E. propinqua</i> ± <i>E. microcorys</i> , <i>Lophostemon confertus</i> , <i>Corymbia intermedia</i> , <i>E. acmenoides</i> open forest on metamorphics ± interbedded volcanics	12.11.3
<i>Corymbia citriodora</i> subsp. <i>variegata</i> , <i>Eucalyptus siderophloia</i> , <i>E. major</i> open forest on metamorphics ± interbedded volcanics	12.11.5
<i>Corymbia citriodora</i> subsp. <i>variegata</i> , <i>Eucalyptus crebra</i> woodland on metamorphics ± interbedded volcanics	12.11.6
<i>Eucalyptus crebra</i> woodland on metamorphics ± interbedded volcanics	12.11.7
Notophyll vine forest ± <i>Araucaria cunninghamii</i> on metamorphics ± interbedded volcanics	12.11.10
Araucarian microphyll vine forest on metamorphics ± interbedded volcanics, usually in southern half of bioregion	12.11.11
<i>Eucalyptus moluccana</i> woodland on metamorphics ± interbedded volcanics	12.11.18
<i>Angophora leiocarpa</i> , <i>Eucalyptus crebra</i> woodland on metamorphics ± interbedded volcanics	12.11.22
<i>Eucalyptus pilularis</i> tall open forest on Mesozoic to Proterozoic igneous rocks especially granite	12.12.2

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Open forest complex with <i>Corymbia citriodora</i> subsp. <i>variegata</i> , <i>Eucalyptus siderophloia</i> or <i>E. crebra</i> or <i>E. decolor</i> , <i>E. major</i> and/or <i>E. longirostrata</i> , <i>E. acmenoides</i> or <i>E. portuensis</i> on Mesozoic to Proterozoic igneous rocks	12.12.3
<i>Eucalyptus acmenoides</i> ± <i>Syncarpia glomulifera</i> tall open forest on Mesozoic to Proterozoic igneous rocks, especially granite	12.12.4
<i>Corymbia citriodora</i> subsp. <i>variegata</i> , <i>Eucalyptus crebra</i> open forest on Mesozoic to Proterozoic igneous rocks	12.12.5
<i>Eucalyptus crebra</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.7
<i>Eucalyptus portuensis</i> or <i>E. acmenoides</i> , <i>Corymbia trachyphloia</i> subsp. <i>trachyphloia</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.11
Araucarian complex microphyll to notophyll vine forest on Mesozoic to Proterozoic igneous rocks	12.12.13
<i>Corymbia intermedia</i> ± <i>Eucalyptus propinqua</i> , <i>E. siderophloia</i> , <i>E. microcorys</i> , <i>Lophostemon confertus</i> open forest on Mesozoic to Proterozoic igneous rocks	12.12.15
Notophyll vine forest on Mesozoic to Proterozoic igneous rocks	12.12.16
<i>Eucalyptus tereticornis</i> ± <i>E. eugenoides</i> woodland on crests, upper slopes and elevated valleys and plains on Mesozoic to Proterozoic igneous rocks	12.12.23

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Angophora leiocarpa</i> , <i>Eucalyptus crebra</i> woodland on Mesozoic to Proterozoic igneous rocks	12.12.24
<i>Eucalyptus fibrosa</i> subsp. <i>fibrosa</i> woodland to open forest on Mesozoic to Proterozoic igneous rocks	12.12.25

Part 13 Wet Tropics Bioregion

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Mangrove closed shrub to open forest of areas subject to regular tidal inundation	7.1.1
<i>Melaleuca quinquenervia</i> and/or <i>Melaleuca cajaputi</i> closed forest to shrubland on poorly drained alluvial plains	7.3.5
<i>Melaleuca viridiflora</i> ± <i>Eucalyptus</i> spp. ± <i>Lophostemon suaveolens</i> open forest to open woodland on poorly drained alluvial plains	7.3.8
<i>Eucalyptus platyphylla</i> woodland to open forest on alluvial plains	7.3.16
<i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> ± <i>E. drepanophylla</i> open forest to open woodland on alluvial plains	7.3.45
Complex mesophyll to mesophyll vine forest on well-drained basalt lowlands and foothills	7.8.1

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Complex mesophyll vine forest of high rainfall, cloudy uplands on basalt, including small areas of wind-sheared notophyll vine forest on ridgelines	7.8.2
Simple to complex notophyll vine forest of cloudy wet highlands on basalt	7.8.4
Simple to complex mesophyll to notophyll vine forest on moderately to poorly drained metamorphics (excluding amphibolites) of moderate fertility of the moist and wet lowlands, foothills and uplands	7.11.1
<i>Eucalyptus pellita</i> ± <i>Corymbia intermedia</i> open forest (or vine forest with <i>E. pellita</i> and <i>C. intermedia</i> emergents) on lowlands and foothills on metamorphics	7.11.5
Complex notophyll vine forest with <i>Agathis robusta</i> emergents on foothills and uplands on metamorphics	7.11.7
Simple notophyll vine forest of moist to very wet metamorphic uplands and highlands	7.11.12
<i>Corymbia nesophila</i> , <i>Corymbia clarksoniana</i> , <i>Eucalyptus platyphyllea</i> open woodland to open forest on gently sloping metamorphic lowlands and foothills	7.11.20
<i>Eucalyptus leptophleba</i> woodland to open forest on metamorphic uplands of the dry rainfall zone	7.11.21
<i>Eucalyptus portuensis</i> ± <i>Corymbia citriodora</i> woodland to open forest on metamorphics	7.11.35
<i>Corymbia clarksoniana</i> and/or <i>Eucalyptus drepanophylla</i> open forest to woodland on metamorphics	7.11.51

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
Simple to complex mesophyll to notophyll vine forest of moderately to poorly drained granites and rhyolites of moderate fertility of the moist and wet lowlands, foothills and uplands	7.12.1
Simple to complex microphyll to notophyll vine forest, often with <i>Agathis robusta</i> or <i>A. microstachya</i> , on granites and rhyolites of moist foothills and uplands	7.12.7
Simple to complex notophyll vine forest and semi-evergreen notophyll vine forest of rocky areas and talus on moist foothills and uplands on granites and rhyolites	7.12.11
Simple to complex notophyll vine forest, including small areas of <i>Araucaria bidwillii</i> , of cloudy wet and moist uplands and highlands on granites and rhyolites	7.12.16
Simple microphyll vine–fern forest with <i>Balanops australiana</i> , <i>Elaeocarpus</i> spp. ± <i>Trochocarpa bellendenkerensis</i> ± <i>Uromyrtus</i> spp. ± <i>Agathis atropurpurea</i> of cloudy wet highlands on granites and rhyolites	7.12.19
<i>Eucalyptus grandis</i> open forest to woodland, or <i>Corymbia intermedia</i> , <i>E. pellita</i> and <i>E. grandis</i> open forest to woodland, (or vine forest with these species as emergents) on granites and rhyolites	7.12.21
<i>Eucalyptus resinifera</i> ± <i>E. portuensis</i> ± <i>Syncarpia glomulifera</i> tall open forest to tall woodland (or vine forest with these species as emergents) of granite and rhyolite uplands and highlands	7.12.22
<i>Eucalyptus portuensis</i> and <i>Corymbia intermedia</i> open forest to woodland (or vine forest with <i>E. portuensis</i> and <i>C. intermedia</i> emergents) on foothills and uplands on granites and rhyolites	7.12.24

[s 7]

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Syncarpia glomulifera</i> ± <i>Corymbia intermedia</i> ± <i>Allocasuarina</i> spp. open forest, or <i>Lophostemon suaveolens</i> , <i>Allocasuarina littoralis</i> , <i>C. intermedia</i> shrubland ± vine forest spp. on exposed ridgelines or steep slopes on granites and rhyolites	7.12.26
<i>Eucalyptus reducta</i> open forest to woodland on uplands and highlands on shallow granitic and rhyolitic soils	7.12.27
<i>Eucalyptus platyphylla</i> ± <i>E. drepanophylla</i> ± <i>Corymbia</i> spp. open woodland to open forest on granites and rhyolites	7.12.28
<i>Corymbia intermedia</i> and/or <i>Lophostemon suaveolens</i> open forest to woodland ± areas of <i>Allocasuarina littoralis</i> and <i>A. torulosa</i> on uplands on granites and rhyolites	7.12.29
<i>Corymbia citriodora</i> ± <i>Eucalyptus portuensis</i> woodland to open forest on granites and rhyolites	7.12.30
<i>Eucalyptus portuensis</i> and/or <i>E. drepanophylla</i> ± <i>C. intermedia</i> ± <i>C. citriodora</i> ± <i>E. granitica</i> open woodland to open forest on uplands on granites	7.12.34
<i>Corymbia clarksoniana</i> ± <i>C. tessellaris</i> ± <i>Eucalyptus drepanophylla</i> ± <i>C. intermedia</i> open forest to woodland, or <i>E. drepanophylla</i> woodland, of moist to dry lowlands, foothills and uplands on granites and rhyolites	7.12.53

Column 1 Regional ecosystem	Column 2 Regional ecosystem number
<i>Eucalyptus tereticornis</i> ± <i>E. granitica</i> woodland to open forest of foothills and uplands on granites and rhyolites	7.12.61
Rock pavements or areas of skeletal soil on granite and rhyolite of dry western or southern areas ± shrublands to closed forests of <i>Acacia</i> spp. and/or <i>Lophostemon suaveolens</i> and/or <i>Allocasuarina littoralis</i> and/or <i>Eucalyptus lockyeri</i> subsp. <i>exuta</i>	7.12.65

8 Amendment of sch 4 (Grassland regional ecosystems—Act, schedule)

Schedule 4, part 11—

insert—

Mixed closed-tussock grassland to closed hermland on coral, shingle and sand cays 12.2.17

9 Amendment of sch 7 (Fees)

(1) Schedule 7, items 7 and 8—

omit.

(2) Schedule 7, item 9—

renumber as schedule 7, item 7.

ENDNOTES

- 1 Made by the Governor in Council on 28 November 2013.
- 2 Notified on the Queensland legislation website on 29 November 2013.
- 3 The administering agency is the Department of Natural Resources and Mines.

© State of Queensland 2013

Authorised by the Parliamentary Counsel